This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

4.1732"
106mm

1/8"R
3.2mm

1.811"
46mm

NDC 80610-430-93

ZORYVE®
(roflumilast)
topical foam, 0.3%

Rx Only | 25 grams

For Topical Use Only

Professional Sample
Not for Sale

Recommended Dosage: See prescribing information.
Not for oral, ophthalmic, or intravaginal use
Store upright. Shake before use. Turn can upside down to dispense.
Keep out of reach of children.
Contains propellant (butane, isobutane, and propane). Store at 20°C to 25°C (68°F to 77°F). Do not freeze.
Warning: Flammable. Avoid fire, flame, and smoking during and immediately following application.
Contents under pressure. Do not puncture or incinerate. Do not expose to heat or store at temperatures above 49°C (120°F).
Manufactured for:
Arcutis Biotherapeutics, Inc.
Westlake Village, CA 91361
000000

ARCUTIS®

000000

3 80610 43093 1

FPO

(b) (4)

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

Lot Number and Expiration Date on Zoryve® Foam Cans

The lot number and expiration date will be printed on the bottom of Zoryve® foam cans as shown below:

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

Each gram contains:
3 mg (0.3%) roflumilast in a white to off-white foam base containing ceteareth-10 phosphate, cetearyl phosphate, cetostearyl alcohol, diethylene glycol monoethyl ether, hexylene glycol, isopropyl palmitate, methylparaben, propylparaben, purified water, sodium hydroxide, and white petrolatum. Hydrochloric acid may have been added to adjust pH. Contains propellant (butane, isobutane, and propane).

Manufactured for:
Arcutis Biotherapeutics, Inc.
Westlake Village, CA 91361

ARCUTIS®

©2023 Arcutis Biotherapeutics, Inc.
All rights reserved.

000000

FPO

3 80610 43093 1

NDC 80610-430-93

ZORYVE®
(roflumilast)
topical foam, 0.3%

Professional Sample
Not for Sale

Rx Only | 25 grams

For Topical Use Only

Recommended Dosage:
See prescribing information.

Not for oral, ophthalmic, or intravaginal use

Store upright. Shake before use. Turn can upside down to dispense.

Keep out of reach of children.

Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F) [See USP Controlled Room Temperature]. Do not freeze.

Warning: Flammable. Avoid fire, flame, and smoking during and immediately following application. Contents under pressure. Do not puncture or incinerate. Do not expose to heat or store at temperatures above 49°C (120°F).

NDC 80610-430-93

ZORYVE®
(roflumilast)
topical foam, 0.3%

Professional Sample
Not for Sale

Rx Only | 25 grams

For Topical Use Only

LOT AAAA-A
EXP YYYY-MMM-DD

000000

(b) (4)

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

David Lewis

Digitally signed by David Lewis
Date: 6/12/2024 03:14:35PM
GUID: 508da72000029f287fa31e664741b577