

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use INLEXZO safely and effectively. See full prescribing information for INLEXZO.

INLEXZO™ (gemcitabine intravesical system)

Initial U.S. Approval: 2025

INDICATIONS AND USAGE

INLEXZO is a nucleoside metabolic inhibitor-containing intravesical system, indicated for the treatment of adult patients with Bacillus Calmette-Guérin (BCG)-unresponsive, non-muscle invasive bladder cancer (NMIBC) with carcinoma *in situ* (CIS) with or without papillary tumors. (1)

DOSAGE AND ADMINISTRATION

For Intravesical Administration Only

- Insert INLEXZO (225 mg of gemcitabine) into the bladder once every 3 weeks up to 6 months (8 doses), followed by once every 12 weeks (6 doses). (2.2)
- Insert into the bladder using the co-packaged urinary catheter and stylet only. (2.1)
- Remove INLEXZO after each 3-week indwelling period. (2.2)
- See Full Prescribing Information and Instructions for Use for insertion and removal procedures. (2.3)

DOSAGE FORMS AND STRENGTHS

- One single-dose 225 mg gemcitabine intravesical system (3)

CONTRAINDICATIONS

- Perforation of the bladder (4, 5.1)
- Prior hypersensitivity reaction to gemcitabine or any component of the product (4)

WARNINGS AND PRECAUTIONS

- **Risks in Patients with Perforated Bladder:** Evaluate the bladder before the intravesical insertion of INLEXZO. Do not administer to patients with a perforated bladder or in whom the integrity of the bladder mucosa has been compromised. (4, 5.1)
- **Risk of Metastatic Bladder Cancer with Delayed Cystectomy:** Delaying cystectomy can lead to the development of metastatic bladder cancer, which can be lethal. (5.2)
- **Magnetic Resonance Imaging (MRI) Safety:** INLEXZO can only be safely scanned with MRI under certain conditions. (5.3)
- **Embryo-Fetal Toxicity:** Can cause fetal harm. Advise patients of the potential risk to a fetus and to use effective contraception. (5.4, 8.1, 8.3)

ADVERSE REACTIONS

- The most common (>15%) adverse reactions, including laboratory abnormalities, are urinary frequency, urinary tract infection, dysuria, micturition urgency, decreased hemoglobin, increased lipase, urinary tract pain, decreased lymphocytes, hematuria, increased creatinine, increased potassium, increased AST, decreased sodium, bladder irritation, and increased ALT. (6.1)

USE IN SPECIFIC POPULATIONS

- **Lactation:** Advise not to breastfeed. (8.2)

To report SUSPECTED ADVERSE REACTIONS, contact Janssen Biotech, Inc. at 1-800-526-7736 (1-800-JANSSEN) or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Issued: 09/2025

FULL PRESCRIBING INFORMATION: CONTENTS*

1	INDICATIONS AND USAGE	
2	DOSAGE AND ADMINISTRATION	
2.1	Important Administration Instructions	
2.2	Recommended Dosage	
2.3	Preparation and Intravesical Administration	
3	DOSAGE FORMS AND STRENGTHS	
4	CONTRAINDICATIONS	
5	WARNINGS AND PRECAUTIONS	
5.1	Risks in Patients with Perforated Bladder	
5.2	Risk of Metastatic Bladder Cancer with Delayed Cystectomy	
5.3	Magnetic Resonance Imaging (MRI) Safety	
5.4	Embryo-Fetal Toxicity	
6	ADVERSE REACTIONS	
6.1	Clinical Trials Experience	
8	USE IN SPECIFIC POPULATIONS	
8.1	Pregnancy	
8.2	Lactation	
8.3	Females and Males of Reproductive Potential	
8.4	Pediatric Use	
8.5	Geriatric Use	
11	DESCRIPTION	
12	CLINICAL PHARMACOLOGY	
12.1	Mechanism of Action	
12.2	Pharmacodynamics	
12.3	Pharmacokinetics	
13	NONCLINICAL TOXICOLOGY	
13.1	Carcinogenesis, Mutagenesis, Impairment of Fertility	
14	CLINICAL STUDIES	
14.1	BCG-unresponsive NMIBC	
15	REFERENCES	
16	HOW SUPPLIED/STORAGE AND HANDLING	
17	PATIENT COUNSELING INFORMATION	

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

INLEXZO is indicated for the treatment of adult patients with Bacillus Calmette-Guérin (BCG)-unresponsive, non-muscle invasive bladder cancer (NMIBC) with carcinoma *in situ* (CIS), with or without papillary tumors.

2 DOSAGE AND ADMINISTRATION

2.1 Important Administration Instructions

Administer INLEXZO intravesically only. Do NOT administer by any other route. INLEXZO is co-packaged with a urinary catheter and stylet used to insert INLEXZO through the urinary catheter into the bladder. Administer using the co-packaged urinary catheter and stylet only.

INLEXZO should be inserted and removed by a trained healthcare provider. Healthcare providers should become thoroughly familiar with the insertion and removal instructions before attempting insertion or removal of INLEXZO.

Prophylactic antibiotics may be used at the discretion of the treating healthcare provider with each INLEXZO insertion and removal.

2.2 Recommended Dosage

Insert INLEXZO (225 mg of gemcitabine) into the bladder once every 3 weeks for up to 6 months (8 doses), followed by once every 12 weeks for up to 18 months (6 doses), or until persistent or recurrent NMIBC, disease progression, or unacceptable toxicity.

Remove INLEXZO after each 3-week indwelling period.

Missed Dose

If a dose is missed, it should be administered as closely as possible to the original treatment schedule.

2.3 Preparation and Intravesical Administration

See the *Instructions for Use* enclosed in the carton for complete information on preparation, intravesical administration, and removal of INLEXZO.

INLEXZO is a hazardous drug. Follow applicable special handling and disposal procedures while handling INLEXZO and during the insertion and removal procedure.¹

Instruct patients to drink approximately 1500 mL of fluids per day during therapy with INLEXZO to ensure adequate urine production for drug release.

Instruct patients not to empty the bladder immediately prior to the insertion procedure. Presence of urine in the bladder can facilitate deployment of INLEXZO. Patients can resume micturition after the insertion procedure.

During indwelling period of approximately 3 weeks, advise patients to avoid urine contact with skin, to void urine sitting on a toilet, to wash hands with soap and water and to wash their genital area with water after each urination, and to flush the toilet after use.

3 DOSAGE FORMS AND STRENGTHS

One single-dose 225 mg strength gemcitabine intravesical system consisting of a flexible bi-oval shaped tube containing an almost white to light pink-brown colored component at the center surrounded by off white to light blue components.

4 CONTRAINDICATIONS

INLEXZO is contraindicated in patients with:

- Perforation of the bladder [*see Warnings and Precautions (5.1)*].
- Prior hypersensitivity reactions to gemcitabine or any component of the product.

5 WARNINGS AND PRECAUTIONS

5.1 Risks in Patients with Perforated Bladder

INLEXZO may lead to systemic exposure to gemcitabine and to severe adverse reactions if administered to patients with a perforated bladder or to those in whom the integrity of the bladder mucosa has been compromised.

Evaluate the bladder before the intravesical administration of INLEXZO and do not administer to patients with a perforated bladder or mucosal compromise until bladder integrity has been restored [*see Contraindications (4)*].

5.2 Risk of Metastatic Bladder Cancer with Delayed Cystectomy

Delaying cystectomy in patients with BCG-unresponsive CIS could lead to development of muscle invasive or metastatic bladder cancer, which can be lethal. The risk of developing muscle invasive or metastatic bladder cancer increases the longer cystectomy is delayed in the presence of persisting CIS.

Of the 83 evaluable patients with BCG-unresponsive CIS treated with INLEXZO in Cohort 2 of SunRISe-1, seven patients (8%) progressed to muscle invasive (T2 or greater) bladder cancer. Three patients (3.5%) had progression determined at the time of cystectomy. The median time between determination of persistent or recurrent CIS or T1 and progression to muscle invasive disease was 94 days.

5.3 Magnetic Resonance Imaging (MRI) Safety

INLEXZO may be used with MRI only under the specific predefined conditions provided below to avoid potential safety hazards or severe adverse reactions.

Based on clinical experience in patients treated with INLEXZO indwelling in the bladder who underwent MRI scans and nonclinical testing, INLEXZO is MR Conditional. A patient with

INLEXZO indwelling in the bladder can be scanned in an MR system under the following conditions:

- Static magnetic field of 1.5-Tesla and 3-Tesla, only.
- Maximum spatial gradient magnetic field of 5000 Gauss/cm or less.
- Maximum magnetic resonance system reported, whole body averaged specific absorption rate of 2-W/kg for 60 minutes of continuous scanning (i.e., per pulse sequence or back-to-back sequences without breaks) in the Normal Operating Mode of operation for the MR system.

Under the scan conditions defined, INLEXZO is expected to produce a maximum temperature rise of 2°C after 15 minutes of continuous scanning.

In nonclinical testing, the image artifact caused by INLEXZO extends approximately 2 mm from INLEXZO using a gradient echo pulse sequence and a 3-Tesla MR system.

5.4 Embryo-Fetal Toxicity

Based on animal data and its mechanism of action, INLEXZO can cause fetal harm when administered to a pregnant woman if systemic exposure occurs [see *Clinical Pharmacology (12.1)*]. In animal reproduction studies, systemic administration of gemcitabine was teratogenic, embryotoxic, and fetotoxic in mice and rabbits.

Advise pregnant women and females of reproductive potential of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment and for 6 months after final removal of INLEXZO. Advise male patients with female partners of reproductive potential to use effective contraception during treatment and for 3 months after final removal of INLEXZO [see *Use in Specific Populations (8.1, 8.3)*].

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The safety of INLEXZO monotherapy was evaluated in Cohort 2 of SunRISe-1, a multi-center, open-label study in 85 adult patients with BCG-unresponsive NMIBC with CIS, with or without papillary tumors [see *Clinical Studies (14.1)*].

Patients received INLEXZO (225 mg of gemcitabine) inserted into the bladder every 3 weeks for 6 months, followed by once every 12 weeks for up to 18 months, or until unacceptable toxicity, disease persistence, recurrence, or progression [see *Dosage and Administration (2.2)*].

The median number of doses of INLEXZO administered to patients was 9 (range: 1 to 14) doses. The median duration of exposure to INLEXZO was 41 weeks (range: 1 to 108 weeks).

Serious adverse reactions occurred in 24% of patients receiving INLEXZO. Serious adverse reactions that occurred in >2% of patients included urinary tract infection, hematuria, pneumonia, and urinary tract pain. Fatal adverse reactions occurred in 1.2% of patients who received INLEXZO, including cognitive disorder.

Permanent discontinuation of INLEXZO due to an adverse reaction occurred in 7% of patients. Adverse reactions which resulted in permanent discontinuation of INLEXZO in >1% of patients included bladder irritation, urinary frequency, cognitive disorder, hydronephrosis, and urinary tract disorder.

Dosage interruptions of INLEXZO due to an adverse reaction occurred in 41% of patients. Adverse reactions which required dosage interruption in >3% of patients included urinary tract infection, urinary tract pain, hematuria, urinary frequency, micturition urgency, dysuria, and genital pain.

The most common (>15%) adverse reactions, including laboratory abnormalities, were urinary frequency, urinary tract infection, dysuria, micturition urgency, decreased hemoglobin, increased lipase, urinary tract pain, decreased lymphocytes, hematuria, increased creatinine, increased potassium, increased AST, decreased sodium, bladder irritation, and increased ALT.

Table 1 summarizes the adverse reactions in SunRISe-1.

Table 1: Adverse Reactions Occurring in >15% of Patients in SunRISe-1

Adverse Reaction	INLEXZO N=85	
	All Grades %	Grade 3 or 4 %
Urinary frequency	48	0
Urinary tract infection ¹	44	6
Dysuria	42	0
Micturition urgency ¹	34	0
Urinary tract pain ¹	26	7
Hematuria ¹	24	2.4
Bladder irritation ¹	16	0

¹ Includes other related terms

Other clinically significant adverse reactions (<15%) included fatigue (14%), genital pain (12%), diarrhea (11%), urinary incontinence (9%), urinary retention (7%), and nocturia (4.7%).

Table 2: Select Laboratory Abnormalities (>15%) That Worsened from Baseline in Patients Who Received INLEXZO in SunRISe-1

Laboratory Abnormality	INLEXZO ¹	
	All Grades (%)	Grade 3 or 4 (%)
Hematology		
Decreased Hemoglobin	31	1.2
Decreased Lymphocytes	24	4.8
Chemistry		
Increased Lipase	28	12
Increased Creatinine	24	0

Increased Potassium	22	1.2
Increased AST	17	1.2
Decreased Sodium	16	4.8
Increased ALT	16	1.2

¹ The denominator used to calculate the rate varied from 82 to 83 based on the number of patients with a baseline value and at least one post-treatment value.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Based on animal data and its mechanism of action, INLEXZO can cause fetal harm when administered to a pregnant woman if systemic exposure occurs [see *Clinical Pharmacology (12.1)*]. There are no available data on the use of INLEXZO in pregnant women to inform a drug-associated risk. In animal reproduction studies, systemic administration of gemcitabine was teratogenic, embryotoxic, and fetotoxic in mice and rabbits (*see Data*). Advise pregnant women and females of reproductive potential of the potential risk to a fetus.

In the U.S. general population, the estimated background risk of major birth defects and miscarriages in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

Data

Animal Data

Gemcitabine is embryotoxic in mice. Daily systemic dosing of gemcitabine to pregnant mice increased the incidence of fetal malformation (cleft palate, incomplete ossification) at doses of 1.5 mg/kg/day. Gemcitabine was embryotoxic and fetotoxic in rabbits. Daily systemic dosing of gemcitabine to pregnant rabbits resulted in fetotoxicity (decreased fetal viability, reduced litter sizes, and developmental delays) and increased the incidence of fetal malformations (fused pulmonary artery, absence of gall bladder) at doses of 0.1 mg/kg/day.

8.2 Lactation

Risk Summary

There is no information regarding the presence of gemcitabine or its metabolites in human milk, or their effects on the breastfed infant or milk production following INLEXZO administration. Because of the potential for serious adverse reactions in breastfed infants, advise women not to breastfeed during treatment and for 1 week after final removal of INLEXZO.

8.3 Females and Males of Reproductive Potential

INLEXZO can cause fetal harm when administered to a pregnant woman [see *Use in Specific Populations (8.1)*].

Pregnancy Testing

Verify pregnancy status in females of reproductive potential prior to initiating INLEXZO.

Contraception

Females

Advise females of reproductive potential to use effective contraception during treatment and for 6 months after final removal of INLEXZO.

Males

Because of the potential for genotoxicity, advise males with female partners of reproductive potential to use effective contraception during treatment and for 3 months after final removal of INLEXZO [see *Nonclinical Toxicology (13.1)*].

Infertility

Males

Based on animal studies, INLEXZO may impair fertility in males of reproductive potential [see *Nonclinical Toxicology (13.1)*]. It is not known whether these effects on fertility are reversible.

8.4 Pediatric Use


Safety and effectiveness of INLEXZO in pediatric patients have not been established.

8.5 Geriatric Use

Of the patients given INLEXZO monotherapy in Cohort 2 of SunRISe-1, 72% were 65 years of age or older and 34% were 75 years or older. There were insufficient numbers of patients <65 years of age to determine if these patients respond differently to patients 65 years of age and older.

11 DESCRIPTION

INLEXZO contains gemcitabine hydrochloride, a nucleoside metabolic inhibitor. Gemcitabine hydrochloride is 2'-deoxy-2',2'-difluorocytidine monohydrochloride (β -isomer) with a molecular formula of $C_9H_{11}F_2N_3O_4 \cdot HCl$, and a molecular weight of 299.66. The structural formula is:



INLEXZO is a sterile, non-resorbable intravesical system containing the equivalent of 225 mg gemcitabine (present as 256.3 mg of gemcitabine hydrochloride).

Gemcitabine Intravesical System

INLEXZO is a bi-oval-shaped tube containing an almost white to light pink-brown colored gemcitabine component at the center surrounded on each side by off white to light blue-colored osmotic components.

- The gemcitabine component contains 225 mg of gemcitabine and the following inactive ingredients: polyethylene glycol 8000 (8.0 mg), povidone K30 (13.4 mg), and urea (42.6 mg).
- The osmotic components contain the following inactive ingredients: FD&C Blue No.1 (0.0042 mg), polyethylene oxide 600,000 (72.0 mg), and urea (648.0 mg).

The silicone tube contains two lumens, the larger one containing the drug components and silicone spacers, and the smaller one containing a superelastic nitinol wire in a predefined shape (wireform). Both lumens are capped with a silicone adhesive. The lumen containing the gemcitabine and osmotic components has a single delivery orifice. INLEXZO's coiled dimensions are approximately 5.5 cm wide × 4.5 cm high.

Urinary Catheter and Stylet

INLEXZO is co-packaged with a sterile urinary catheter and a sterile stylet, required for transurethral insertion into the bladder. The urinary catheter and stylet are made of thermoplastic elastomer and polyethylene, and consist of the following components:

- A semi-transparent urinary catheter, with a rounded blunt distal tip that includes a coude tip, a product exit port near the distal tip, and a lumen which extends from the exit port to an open proximal end. The outer diameter of the urinary catheter is 5.82 mm (17.5 Fr). Printed depth markings are placed on the urinary catheter to indicate insertion depth and orientation of the coude tip to assist the user during insertion.
- A green stylet with a hub at the proximal end is used to advance INLEXZO through the urinary catheter lumen and into the bladder. The stylet length and proximal hub prevent the stylet's distal end from advancement beyond the exit port.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Gemcitabine kills cells undergoing DNA synthesis and blocks the progression of cells through the G1/S-phase boundary. Gemcitabine is metabolized by nucleoside kinases to diphosphate (dFdCDP) and triphosphate (dFdCTP) nucleosides. Gemcitabine diphosphate inhibits ribonucleotide reductase, an enzyme responsible for catalyzing the reactions that generate deoxynucleoside triphosphates for DNA synthesis, resulting in reductions in deoxynucleotide concentrations, including dCTP. Gemcitabine triphosphate competes with dCTP for incorporation into DNA. The reduction in the intracellular concentration of dCTP by the action of the diphosphate enhances the incorporation of gemcitabine triphosphate into DNA (self-potential). After the gemcitabine nucleotide is incorporated into DNA, only one additional nucleotide is added to the growing DNA strands, which eventually results in the initiation of apoptotic cell death.

12.2 Pharmacodynamics

The exposure-response relationship and time-course of pharmacodynamic response for the safety and effectiveness of INLEXZO have not been fully characterized.

12.3 Pharmacokinetics

The systemic exposure of gemcitabine and the inactive uracil metabolite (2'-deoxy-2',2'-difluorouridine [dFdU]), were evaluated between Days 2 and 7 during the indwelling period. The plasma gemcitabine concentrations were below the lower limit of quantification (0.1 µg/mL) in all patients at all time points. Eighteen patients (17%) had at least one quantifiable plasma dFdU concentration above the lower limit of quantification (0.1 µg/mL) and the maximum observed plasma dFdU concentration was 0.4 µg/mL. Plasma concentrations of both gemcitabine and dFdU are estimated to be less than 1% of the expected C_{max} after intravenous administration of gemcitabine.

Excretion

Gemcitabine and dFdU are excreted in urine throughout the indwelling period for INLEXZO. Of the total gemcitabine dose, 77% was excreted by Day 7 and 99% was excreted by Day 21 in urine as gemcitabine and dFdU.

Mean urinary excretion on Days 2 through 5 ranged from 21 to 32 mg per day excreted in the urine as gemcitabine and dFdU.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term animal studies to evaluate the carcinogenic potential of gemcitabine have not been conducted. Gemcitabine was mutagenic in an *in vitro* mouse lymphoma (L5178Y) assay and was clastogenic in an *in vivo* mouse micronucleus assay.

Dedicated animal fertility studies have not been conducted with gemcitabine intravesical system. Gemcitabine intraperitoneal doses of 0.5 mg/kg/day in male mice resulted in moderate to severe hypospermatogenesis, decreased fertility, and decreased implantations. In female mice, fertility was not affected but maternal toxicities were observed at 1.5 mg/kg/day administered intravenously and fetotoxicity or embryo lethality were observed at 0.25 mg/kg/day administered intravenously.

14 CLINICAL STUDIES

14.1 BCG-unresponsive NMIBC

The efficacy of INLEXZO was evaluated in Cohort 2 of SunRISe-1 (NCT04640623), a single-arm, multi-center trial in 83 adults with BCG-unresponsive, NMIBC with CIS, with or without papillary tumors (T1, or high-grade Ta) following transurethral resection.

BCG-unresponsive NMIBC CIS was defined as persistent or recurrent CIS alone or with Ta/T1 disease within 12 months of adequate BCG therapy. Adequate BCG therapy was defined as a

minimum administration of at least five of six doses of an initial induction course plus either of: at least two of three doses of maintenance therapy or at least two of six doses of a second induction course. Prior to treatment, all patients had undergone transurethral resection of bladder tumor (TURBT) to remove all resectable disease (Ta and T1 components). Residual CIS (Tis components) not amenable to complete resection was permitted. The trial included patients who were ineligible for or who had elected not to undergo radical cystectomy and excluded patients with extra-vesical (i.e., urethra, ureter, or renal pelvis), muscle invasive (T2-T4), locally advanced, or metastatic urothelial carcinoma.

Patients received INLEXZO (225 mg of gemcitabine) into the bladder every 3 weeks for 6 months, followed by once every 12 weeks for up to 18 months, or until unacceptable toxicity, persistence or recurrence of CIS and/or high-grade papillary disease, or progression [see *Dosage and Administration (2.2)*]. Tumor status was assessed every 12 weeks during the initial two years of treatment, after which cystoscopy was performed at least every 24 weeks. Mandatory biopsies were performed 24 and 48 weeks after treatment initiation.

The major efficacy outcome measures were complete response rate at any time (defined as negative results for cystoscopy [with TURBT and centrally-reviewed biopsies as applicable] and centrally-reviewed urine cytology) and duration of response.

The median age of patients was 71 years (range: 40 to 88); 80% male; 87% White, 10% Asian, 2.4% Black or African American and 1.2% race not reported; 10% were of Hispanic or Latino ethnicity, 89% identified as not Hispanic or Latino, and 1.2% were ethnicity unknown or not reported. Baseline ECOG performance status was 0 (92%) or 1 (8%). Tumor pattern at study entry was CIS only (67%), CIS with high-grade Ta only (22%), and CIS with T1 (11%). All patients had predominant urothelial carcinoma including 1 patient (1.2%) with squamous differentiation. The median number of prior instillations of BCG was 12 (range: 7 to 42).

Efficacy results are summarized in Table 3.

Table 3: Efficacy Results in SunRISe-1

Endpoint	INLEXZO N=83
Complete Response Rate (95% CI)	82% (72, 90)
Duration of Response^a	
Range in months	0+, 44+
% (n) with duration ≥12 months	51% (35)

CI= confidence interval.

^a Based on patients (n=68) with a complete response at any time; reflects period from the time complete response was achieved.

+ Denotes ongoing response

15 REFERENCES

1. OSHA Hazardous Drugs. Occupational Safety and Health Administration. Available at: <https://www.osha.gov/hazardous-drugs> (Accessed: 06 September 2024).

16 HOW SUPPLIED/STORAGE AND HANDLING

How Supplied

INLEXZO (gemcitabine intravesical system) contains 225 mg gemcitabine.

INLEXZO (NDC# 57894-225-01) is available in a carton containing:

- One sterile single-dose of INLEXZO in two clear laminate sleeves and packaged in an inner pouch. The inner pouch and a desiccant are packaged in an outer foil pouch. The outside surfaces of the inner and outer pouches are not sterile.
- One sterile urinary catheter and one sterile stylet packaged together in a pouch.

Storage

Store in the original carton at 20°C to 25°C (68°F to 77°F); with excursions permitted between 15°C to 30°C (59°F to 86°F) [see *USP Controlled Room Temperature*].

Handling

INLEXZO is a hazardous drug. Follow applicable special handling and disposal procedures.¹

17 PATIENT COUNSELING INFORMATION

Advise patients to read the FDA-approved patient labeling (Patient Information).

Risk of Metastatic Bladder Cancer with Delayed Cystectomy

- Inform patients that delaying cystectomy could lead to development of metastatic bladder cancer. Discuss the risk of metastatic bladder cancer and that the risk increases the longer cystectomy is delayed in the presence of persistent CIS [see *Warnings and Precautions (5.2)*].

Magnetic Resonance Imaging (MRI) Safety

- Inform patients that INLEXZO can only be safely scanned with MRI under specific conditions [see *Warnings and Precautions (5.3)*]. Instruct patients who will have an MRI to tell their healthcare provider that they have INLEXZO.
- This information is included in the MRI Safety Information Card. Complete the MRI Safety Information Card and give it to the patient.

Embryo-Fetal Toxicity

- Advise females of reproductive potential of the potential risk to a fetus and to inform their healthcare provider of a known or suspected pregnancy [see *Warnings and Precautions (5.4) and Use in Specific Populations (8.1)*].
- Advise females of reproductive potential to use effective contraception during treatment and for 6 months after final removal of INLEXZO [see *Use in Specific Populations (8.3)*].

- Advise males with female partners of reproductive potential to use effective contraception during treatment and for 3 months after final removal of INLEXZO [see *Use in Specific Populations (8.3)*].

Lactation

- Advise women not to breastfeed during treatment and for 1 week after final removal of INLEXZO [see *Use in Specific Populations (8.2)*].

Infertility

- Advise males of reproductive potential that INLEXZO may impair fertility [see *Use in Specific Populations (8.3)*].

Important Post-Treatment Instructions

- Instruct patients not to empty the bladder immediately prior to the insertion procedure.
- Advise patients to avoid contact with urine while INLEXZO is indwelling in the bladder and for at least 24 hours post-removal [see *Clinical Pharmacology (12.3)*].
- Advise patients to avoid urine contact with skin by voiding sitting on a toilet, flushing the toilet after use, and to wash hands with soap and water and to wash their genital area with water after each urination.
- Advise patients to wash clothing soiled with urine promptly and separately from other clothing [see *Dosage and Administration (2.3)*].

Manufactured for:
Janssen Biotech, Inc.
Horsham, PA 19044, USA
U.S. License Number 1864

For patent information: www.janssenpatents.com

© Johnson & Johnson and its affiliates 2025

PATIENT INFORMATION
INLEXZO™ (inn-lex-zo)
(gemcitabine intravesical system)

What is INLEXZO?

- INLEXZO is a prescription medicine for the treatment of adults with a type of cancer of the lining of the bladder called non-muscle invasive bladder cancer (NMIBC), that has not spread to other parts of the body, and that did not respond to treatment with Bacillus Calmette-Guérin (BCG). It is not known if INLEXZO is safe and effective for use in children.

Do not receive INLEXZO if you:

- have a tear or hole (perforation) of your bladder.
- have had an allergic reaction to gemcitabine or any of the ingredients in INLEXZO. See the end of this Patient Information leaflet for a complete list of ingredients in INLEXZO.

Before receiving INLEXZO, tell your healthcare provider about all of your medical conditions, including if you:

- are pregnant, or plan to become pregnant. INLEXZO can harm your unborn baby. You should not become pregnant during treatment with INLEXZO. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with INLEXZO.

Females who are able to become pregnant:

- Your healthcare provider will check to see if you are pregnant before starting treatment with INLEXZO.
- Use effective birth control (contraception) during treatment with INLEXZO and for 6 months after final removal of INLEXZO.

Males treated with INLEXZO:

- If you have a female partner who is able to become pregnant, you should use effective birth control (contraception) during treatment with INLEXZO and for 3 months after final removal of INLEXZO.
- are breastfeeding, or plan to breastfeed. It is not known if INLEXZO passes into your breastmilk. Do not breastfeed during treatment with INLEXZO and for 1 week after final removal of INLEXZO.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

How will I receive INLEXZO?

- INLEXZO will be inserted and removed by your healthcare provider.
- INLEXZO is inserted into your bladder through a tube called a urinary catheter 1 time every 3 weeks for up to 6 months (8 doses) and then 1 time every 12 weeks for up to 18 months (6 doses).
- INLEXZO is removed after 3 weeks of being inserted (3-week indwelling period).
- Your healthcare provider will decide how many INLEXZO treatments you will receive.
- Your healthcare provider may give you antibiotics before INLEXZO is inserted or removed.
- It is very important that you go to all of your appointments. If you miss any appointments, call your healthcare provider as soon as possible to schedule your appointment.

Before receiving INLEXZO:

- Do not empty your bladder right before your procedure to insert INLEXZO.

After receiving INLEXZO:

- Drink about 6 to 7 cups (1500 mL) of fluids per day during treatment with INLEXZO to make sure you produce enough urine for the medicine to be released into the bladder.
- You can urinate normally. There is no need to hold your urine.
- Avoid contact between your skin and urine.
- To urinate, **males and females** should sit on the toilet and flush after each use.
- Wash your hands with soap and water and wash your genital area with water after each time you urinate.
- Wash clothing soiled with urine right away and separately from other clothing.

After your healthcare provider removes INLEXZO:

- Avoid contact between your skin and urine for at least 24 hours after INLEXZO is removed.

What should I avoid after INLEXZO is inserted?

Tell your healthcare provider that you have INLEXZO before having a type of scan called Magnetic Resonance Imaging (MRI).

INLEXZO may be used under specific conditions. Your healthcare provider will give you an MRI Safety Information Card. Keep the card in a safe place and show it to all of your healthcare providers. The card contains important information in case you need to have an MRI. Your healthcare provider will review the information on the MRI Safety Information Card and determine the conditions they can safely do an MRI scan while INLEXZO is in your bladder.

What are the possible side effects of INLEXZO?

The most common side effects of INLEXZO include:

- | | |
|--|--|
| • frequent need to pass urine more often than usual | • decreased lymphocytes |
| • urinary tract infection | • blood in urine |
| • pain or burning sensation when passing urine | • increased creatinine |
| • urgent need to pass urine | • increased potassium |
| • decreased hemoglobin | • increased aspartate aminotransferase |
| • increased lipase | • decreased sodium |
| • pain in the urinary tract (felt in the lower stomach area or lower back) | • bladder irritation |
| | • increased alanine transaminase |

INLEXZO may cause fertility problems in males, which may affect your ability to have children. It is unknown if these side effects of fertility are reversible. Talk to your healthcare provider if this is a problem for you.

Tell your healthcare provider if you have any side effect that bothers you or does not go away.

These are not all the possible side effects of INLEXZO.

Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

General information about the safe and effective use of INLEXZO.

Medicines are sometimes prescribed for purposes other than those listed in this Patient Information leaflet. You can ask your pharmacist or healthcare provider for information about INLEXZO that is written for healthcare professionals.

What are the ingredients in INLEXZO?

Active ingredient: gemcitabine hydrochloride

Inactive ingredients for gemcitabine component: polyethylene glycol 8000, povidone K30, and urea.

Inactive ingredients for osmotic components: FD&C Blue No.1, polyethylene oxide 600,000, and urea.

Manufactured for:
Janssen Biotech, Inc.
Horsham, PA 19044, USA
U.S. License Number 1864


For patent information: www.janssenpatents.com
© Johnson & Johnson and its affiliates 2025
For more information, call 1-800-526-7736 or go to www.INLEXZO.com

This Patient Information has been approved by the U.S. Food and Drug Administration

Issued: 09/2025

INSTRUCTIONS FOR USE
INLEXZO™
[inn-lex-zo]
(gemcitabine intravesical system)
For Intravesical Administration Only

INLEXZO™
Intravesical System




Intended Use

INLEXZO releases gemcitabine into bladder urine during the indwelling period. See INLEXZO Prescribing Information (PI) for additional details.

The contents of the Instructions for Use provide instructions for both the insertion and removal of INLEXZO. Ensure that the Instructions for Use are available and reviewed prior to insertion and removal of INLEXZO.

User

INLEXZO should be inserted and removed by a trained healthcare provider.

 MR Conditional	<p>MRI Safety Information INLEXZO is MR Conditional. A patient with INLEXZO may be safely scanned under the following conditions. Failure to follow these conditions may result in injury to the patient.</p>	
Parameter	Condition of Use/Information	
Nominal Values of Static Magnetic Field (T)	1.5-Tesla and 3.0-Tesla	

Maximum Spatial Field Gradient (T/m and gauss/cm)	50-T/m (5,000-gauss/cm)
Type of RF Excitation	Circularly Polarized (CP) (i.e., Quadrature Transmission)
Transmit RF Coil	Any transmit RF coil may be used.
Receive RF Coil	Any receive-only RF coil may be used
Maximum Whole Body Averaged SAR (W/kg)	2-W/kg (Normal Operating Mode)
Limits on Scan Duration	Whole body averaged SAR of 2-W/kg for 60 minutes of continuous RF exposure (i.e., per pulse sequence or back-to-back sequences/series without breaks)
MR Image Artifact	The presence of INLEXZO produces an imaging artifact. Therefore, carefully select pulse sequence parameters if INLEXZO is located in the area of interest.

Precautions

INLEXZO is a hazardous drug. Follow applicable special handling and disposal procedures while handling INLEXZO and during the insertion and removal procedure. Dispose of the used urinary catheter and stylet, INLEXZO, and its packaging per facility procedures and per applicable federal, state, and local regulations.

Wear gloves, and take appropriate precautions, per local guidelines for handling hazardous drugs, to prevent skin or mucus membrane exposure while handling INLEXZO and during the insertion and removal procedure.

If contact with INLEXZO is suspected, immediately wash the skin thoroughly or rinse the mucosa with copious amounts of water.

Advise patients and caregivers to **exercise caution when handling urine during indwelling period of approximately 3 weeks. See INLEXZO PI for details.**

INLEXZO contains one single-dose 225 mg strength gemcitabine intravesical system consisting of a flexible bi-oval shaped tube containing an almost white to light pink-brown colored component at the center surrounded by off white to light blue osmotic components.

Important Information

Use aseptic technique during insertion and removal of INLEXZO.

Follow these instructions carefully, to avoid patient injury and ensure proper functioning.

Insertion

To ensure proper insertion of INLEXZO and to avoid damage to INLEXZO, use only the following:

- Water-based lubricant
- Urinary catheter and stylet (supplied)

Do not use the urinary catheter and stylet for any other purpose. Do not re-sterilize/re-use the urinary catheter or stylet. Re-use of the urinary catheter and stylet can lead to its degradation, failure, and contamination, which can increase the risk of infection or transmission of blood borne pathogens to patients and users.

Do not use any components that are damaged or have damaged packaging.

Check the expiration ('EXP') date before use.

Do not use INLEXZO if expiration date has passed.

Removal


To ensure proper INLEXZO removal and to avoid damage to INLEXZO and/or cystoscope, use only the following:

- **Non-cutting**, grasping forceps
- Flexible or rigid cystoscope


Storage

Store in the original package at **20°C to 25°C** (68°F to 77°F); with excursions permitted between **15°C to 30°C** (59°F to 86°F) [see *USP Controlled Room Temperature*].

INLEXZO™ Intravesical System



Urinary Catheter and Stylet



Prepare components for INLEXZO insertion

1. Gather supplies


Included in the product carton:

- One sterile INLEXZO
- One sterile urinary catheter and one sterile stylet

Not included in the product carton:

- Multiple pairs of gloves
 - Two 10 mL prefilled water-based lubricant syringes
- OR**
- Two empty 10 mL syringes and water-based lubricant

2. Put on gloves





3. Prepare two syringes with sterile water-based lubricant

- Remove each sterile syringe prefilled with **water-based** lubricant from its packaging and place each syringe on the sterile work surface using aseptic technique.


OR

- Fill each of the two empty syringes with 2 mL to 3 mL of **water-based** lubricant plus additional lubricant for urinary catheter tip lubrication.
- Place each lubricant syringe on a sterile work surface using aseptic technique.
- The lubricant is to be used to lubricate the tip of the urinary catheter and to facilitate the insertion of INLEXZO.



4. Open the INLEXZO foil pouch

1. Tear open the INLEXZO outer foil pouch at tear notch.
2. Remove the white INLEXZO inner pouch.




5. Examine the white component pouches for damage

Check the white INLEXZO pouch and white pouch containing the urinary catheter and stylet for damage (e.g., cuts, tears, punctures) that could compromise sterility of the components before opening.


⚠ Do not use if packaging is damaged.

⚠ Surfaces of the white INLEXZO pouch and the white component pouches are not sterile.




6. Open the white component pouches and transfer the contents onto a sterile work surface

1. Open the white INLEXZO pouch and transfer INLEXZO onto a sterile work surface.
2. Do not remove plastic sleeves from INLEXZO.
3. Open the white pouch containing the urinary catheter and stylet and transfer the contents onto a sterile work surface.




7. Put on sterile gloves

1. Ensure your patient is prepared for the procedure.
2. Put on sterile gloves.



8. Remove plastic sleeves

Remove the plastic sleeves from INLEXZO.




9. Inspect components

Inspect INLEXZO, the urinary catheter, and green stylet for damage.

Do not use if the urinary catheter or green stylet are damaged or if the outer surface of INLEXZO is damaged.

Insert INLEXZO



1. Lubricate the urinary catheter

Lubricate tip of the urinary catheter.

The urinary catheter should be free of INLEXZO and the stylet.

2. Insert the urinary catheter (without stylet)

Introduce the urinary catheter into the urethra by hand and advance until urine return. Do not empty the bladder.


Use depth markings to maintain coude tip orientation and insertion depth position throughout the procedure.

⚠ Do not excessively force the urinary catheter into the bladder

⚠ In case of resistance, careful assessment and standard procedural techniques can be performed, as appropriate, in order to safely advance the urinary catheter into the bladder.


⚠ If advancement is obstructed or cause cannot be determined or resolved, withdraw the urinary catheter to avoid patient injury or urinary catheter damage.

⚠ Do not re-use urinary catheter.




3. Inject lubricant from the first syringe into the urinary catheter

Inject 2 mL to 3 mL of lubricant from the first syringe into the end of the urinary catheter with the urinary catheter placed in the bladder.




4. Insert INLEXZO into the urinary catheter

Insert either end of INLEXZO into the urinary catheter and advance until fully inserted.



5. Inject lubricant from the second syringe into the urinary catheter

Inject 2 mL to 3 mL of lubricant from the second syringe into the urinary catheter to help advance INLEXZO further, with the urinary catheter placed in the bladder.



6. Insert stylet into the urinary catheter


Slowly insert the stylet into the urinary catheter until the stylet hub is flush with end of the urinary catheter. This ensures INLEXZO exits the urinary catheter and enters the bladder.

⚠ If INLEXZO cannot be advanced, remove the urinary catheter and stylet together as a single unit.

Ensure INLEXZO is also removed.

Do not attempt to re-use the removed INLEXZO.

Begin again by obtaining a new carton of INLEXZO including a new urinary catheter and stylet.




7. Remove the urinary catheter and stylet together as a single unit

Do not remove the urinary catheter and stylet individually.

INLEXZO should remain inside the bladder.

Retain Instructions for Use for the removal procedure.

Post-Insertion Instructions



1. Provide the completed MRI Safety Information Card to the patient

Remove the MRI Safety Information Card from the carton.

Complete the details and give it to the patient.

Advise the patient to carry the card and to show their current and future healthcare providers in case of need for MRI scans.

- INLEXZO contains a metal wire. The patient can safely undergo an MR exam only under very specific conditions (see MRI Safety Information).


2. Inform the patient and caregivers about the indwelling period

The indwelling period is approximately 3 weeks. See INLEXZO PI for additional information.

Inform the patient and caregivers that INLEXZO will remain in the bladder for the indwelling dosing period.

INLEXZO contains a hazardous drug. The patient and caregivers should be made aware of the need to **exercise caution when handling urine during the indwelling period**. See INLEXZO PI for additional information.

Removal of INLEXZO after the indwelling period



1. Lubricate the cystoscope

Use a water-based lubricant to lubricate the cystoscope.

2. Insert the cystoscope

Insert the cystoscope into the bladder to locate INLEXZO.




3. Grasp INLEXZO

Introduce **non-cutting** grasping forceps through the working channel of the cystoscope.


⚠ Do not use cutting forceps.

Grasp INLEXZO over tubing and metal wire.

⚠ Do not grasp on or near the ends of INLEXZO.



Grasping near the ends of INLEXZO could result in exposing the metal wire, potentially causing damage to surrounding tissue.



4. Remove INLEXZO

Remove the cystoscope and forceps out of the urethra **together** to remove INLEXZO under direct vision.

Do not remove INLEXZO through working channel of the cystoscope. Doing so may damage INLEXZO and/or the cystoscope.

	5. Inspect INLEXZO
--	---------------------------

⚠ After removal, inspect INLEXZO to confirm it is intact and unbroken.

Manufactured for:
Janssen Biotech, Inc.
Horsham, PA 19044, USA.
U.S. License Number 1864

This Instructions for Use has been approved by the U.S. Food and Drug Administration.
Approved: 09/2025

For patent information:
www.janssenpatents.com
© Johnson & Johnson and its affiliates 2025