

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use DOPELET safely and effectively. See full prescribing information for DOPELET.

DOPELET® (avatrombopag) tablets, for oral use
DOPELET® SPRINKLE (avatrombopag) oral granules
Initial U.S. Approval: 2018

RECENT MAJOR CHANGES

Indications and Usage (1.3)	07/2025
Dosage and Administration (2.1, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8)	07/2025
Warnings and Precautions (5.1)	07/2024

INDICATIONS AND USAGE

DOPELET is a thrombopoietin receptor agonist indicated for the treatment of:

- Thrombocytopenia in adult patients with chronic liver disease who are scheduled to undergo a procedure. (1.1)
- Thrombocytopenia in adult patients with chronic immune thrombocytopenia who have had an insufficient response to a previous treatment (1.2)
- Thrombocytopenia in pediatric patients 1 year and older with persistent or chronic immune thrombocytopenia who have had an insufficient response to a previous treatment (1.3)

DOSAGE AND ADMINISTRATION

- DOPELET tablets and DOPELET SPRINKLE are not substitutable on a mg-to-mg basis.
- DOPELET SPRINKLE capsules should be opened, and the contents (oral granules) mixed with a soft food or liquid. Administer immediately after mixing. Do not swallow the capsules whole.
- Administer DOPELET tablets and DOPELET SPRINKLE with food. (2.1, 2.3, 2.5)
- **Chronic Liver Disease:** Dose DOPELET tablets based upon platelet count prior to procedure, orally for 5 days beginning 10 days to 13 days before procedure. For platelet count less than $40 \times 10^9/L$, the dose is 60 mg (3 tablets) orally once daily; for platelet count 40 to less than $50 \times 10^9/L$ the dose is 40 mg (2 tablets) orally once daily. (2.2)
- **Adult Patients with Chronic Immune Thrombocytopenia and Pediatric Patients 6 Years and Older with Persistent or Chronic Immune Thrombocytopenia:** Initiate DOPELET tablets at 20 mg (1 tablet) orally once daily. Adjust the dose or frequency of dosing to maintain platelet count greater than or equal to $50 \times 10^9/L$. Do not exceed 40 mg (2 tablets) per day.

- **Pediatric Patients 1 Year to Less than 6 Years with Persistent or Chronic Immune Thrombocytopenia:** Initiate DOPELET SPRINKLE oral granules at 10 mg (content of 1 capsule) orally once daily. Adjust the dose or frequency of dosing to maintain platelet count greater than or equal to $50 \times 10^9/L$. Do not exceed 20 mg (content of 2 capsules) per day.

DOSAGE FORMS AND STRENGTHS

Tablet: 20 mg (3)
Oral Granules: 10 mg (3)

CONTRAINDICATIONS

None. (4)

WARNINGS AND PRECAUTIONS

Thrombotic/Thromboembolic Complications: DOPELET is a thrombopoietin (TPO) receptor agonist and TPO receptor agonists have been associated with thrombotic and thromboembolic complications in patients with chronic liver disease or immune thrombocytopenia.

Monitor platelet counts. Monitor for signs and symptoms of thromboembolic events and institute treatment promptly. (5.1)

ADVERSE REACTIONS

In adult patients with chronic liver disease, the most common adverse reactions ($\geq 3\%$) were pyrexia, abdominal pain, nausea, headache, fatigue, and edema peripheral. (6.1)

In adult patients with chronic immune thrombocytopenia, the most common adverse reactions ($\geq 10\%$) were headache, fatigue, contusion, epistaxis, upper respiratory tract infection, arthralgia, gingival bleeding, petechiae and nasopharyngitis. (6.1)

In pediatric patients with persistent or chronic immune thrombocytopenia, the most common adverse reactions ($\geq 10\%$) were viral infection, nasopharyngitis, cough, pyrexia, and oropharyngeal pain. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Sobi, Inc. at 1-866-773-5274 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

Moderate or Strong Dual CYP2C9 and CYP3A4 Inducers or Inhibitors: Dose adjustments are recommended for patients with persistent or chronic immune thrombocytopenia. (7.1)

USE IN SPECIFIC POPULATIONS

- Pregnancy: May cause fetal harm (8.1)
- Lactation: Advise not to breastfeed. (8.2)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 07/2025

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

- 1.1 Treatment of Thrombocytopenia in Patients with Chronic Liver Disease
- 1.2 Treatment of Thrombocytopenia in Adult Patients with Chronic Immune Thrombocytopenia
- 1.3 Treatment of Thrombocytopenia in Pediatric Patients 1 Year and Older with Persistent or Chronic Immune Thrombocytopenia

2 DOSAGE AND ADMINISTRATION

- 2.1 Important Use and Administration Instructions
- 2.2 Recommended Dosage of DOPELET Tablets for Patients with Chronic Liver Disease
- 2.3 Recommended Dosage of DOPELET Tablets for Adult Patients with Chronic Immune Thrombocytopenia and Pediatric Patients 6 Years and Older with Persistent or Chronic Immune Thrombocytopenia
- 2.4 Recommended Initial Dosage of DOPELET Tablets with Concomitant Moderate or Strong Dual Inducers or Inhibitors of CYP2C9 and CYP3A4

- 2.5 Recommended Dosage of DOPELET SPRINKLE for Patients 1 Year to Less than 6 Years with Persistent or Chronic Immune Thrombocytopenia
- 2.6 Recommended Initial Dosage of DOPELET SPRINKLE with Concomitant Moderate or Strong Dual Inducers or Inhibitors of CYP2C9 and CYP3A4
- 2.7 Important Preparation and Administration Instructions for DOPELET SPRINKLE
- 2.8 Missed Dose

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Thrombotic/Thromboembolic Complications

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

7 DRUG INTERACTIONS

- 7.1 Effect of Other Drugs on DOPTELET in Patients with Persistent or Chronic Immune Thrombocytopenia

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
8.2 Lactation
8.4 Pediatric Use
8.5 Geriatric Use

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
12.2 Pharmacodynamics
12.3 Pharmacokinetics
12.5 Pharmacogenomics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

- 14.1 Patients with Chronic Liver Disease
14.2 Adult Patients with Chronic Immune Thrombocytopenia
14.3 Pediatric Patients with Persistent or Chronic Immune Thrombocytopenia

16 HOW SUPPLIED/STORAGE AND HANDLING

- 16.1 DOPTELET Tablets
16.2 DOPTELET SPRINKLE

17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

1.1 Treatment of Thrombocytopenia in Patients with Chronic Liver Disease (CLD)

DOPTELET is indicated for the treatment of thrombocytopenia in adult patients with chronic liver disease who are scheduled to undergo a procedure.

1.2 Treatment of Thrombocytopenia in Adult Patients with Chronic Immune Thrombocytopenia (ITP)

DOPTELET is indicated for the treatment of thrombocytopenia in adult patients with chronic immune thrombocytopenia who have had an insufficient response to a previous treatment.

1.3 Treatment of Thrombocytopenia in Pediatric Patients 1 Year and Older with Persistent or Chronic Immune Thrombocytopenia (ITP)

DOPTELET is indicated for the treatment of thrombocytopenia in pediatric patients 1 year and older with persistent or chronic immune thrombocytopenia who have had an insufficient response to a previous treatment.

2 DOSAGE AND ADMINISTRATION

2.1 Important Use and Administration Instructions

- Select the recommended product (DOPTELET tablets or DOPTELET SPRINKLE) based on the indication and patient's age.
- Administer DOPTELET tablets and DOPTELET SPRINKLE with food.
- DOPTELET tablets and DOPTELET SPRINKLE are not substitutable on a mg-to-mg basis.
- The mixture prepared from the granules in DOPTELET SPRINKLE capsules is more bioavailable than DOPTELET tablets [see *Clinical Pharmacology (12.3)*]. There is no experience from clinical trials in switching between dosing with the granules and the tablet. If the formulation is switched, monitor platelet counts weekly until stable platelet counts are obtained and adjust dosing as needed before resuming monthly monitoring.

2.2 Recommended Dosage of DOPTOLET Tablets for Patients with Chronic Liver Disease

Begin DOPTOLET tablets dosing 10 days to 13 days prior to the scheduled procedure. The recommended daily dose of DOPTOLET is based on the patient's platelet count prior to the scheduled procedure (see [Table 1](#)). Patients should undergo their procedure 5 days to 8 days after the last dose of DOPTOLET.

DOPTOLET tablets should be taken orally once daily for 5 consecutive days with food. All 5 days of dosing should be completed.

Table 1: Recommended DOPTOLET Tablets Dosage and Duration in Patients with Chronic Liver Disease Scheduled to Undergo a Procedure

Platelet Count	Recommended DOPTOLET Dosage	Duration
Less than $40 \times 10^9/L$	60 mg (3 tablets) orally once daily	5 days
$40 \times 10^9/L$ to less than $50 \times 10^9/L$	40 mg (2 tablets) orally once daily	5 days

DOPTOLET tablets have been investigated only as a single 5-day once daily dosing regimen in clinical trials in patients with chronic liver disease [see *Clinical Studies (14.1)*]. DOPTOLET should not be administered to patients with chronic liver disease in an attempt to normalize platelet counts.

Monitoring: Obtain a platelet count prior to administration of DOPTOLET therapy and on the day of a procedure to ensure an adequate increase in platelet count.

2.3 Recommended Dosage of DOPTOLET Tablets for Adult Patients with Chronic Immune Thrombocytopenia and Pediatric Patients 6 Years and Older with Persistent or Chronic Immune Thrombocytopenia

Use the lowest dose of DOPTOLET needed to achieve and maintain a platelet count greater than or equal to $50 \times 10^9/L$ as necessary to reduce the risk for bleeding. Dose adjustments are based on platelet count response. Do not use DOPTOLET to normalize platelet counts.

Initial Dosage:

- Begin DOPTOLET tablets at an initial dosage of 20 mg (1 tablet) orally once daily with food (see [Table 3](#)).
- The recommended initial dosages of DOPTOLET tablets are different for patients receiving certain concomitant medications (see [Table 4](#)).

Monitoring: After initiating therapy with DOPTOLET tablets, assess platelet counts weekly until a stable platelet count greater than or equal to $50 \times 10^9/L$ has been achieved, and then obtain platelet counts monthly thereafter. Obtain platelet counts weekly for at least 4 weeks following discontinuation of DOPTOLET.

Dose Adjustments:

DOPTOLET tablet dose adjustments (see [Table 2](#) and [Table 3](#)) are based on the platelet count response. Do not exceed a DOPTOLET daily dose of 40 mg (2 tablets).

Table 2: DOPTELET Tablets Recommended Dose Adjustments for Adult Patients with Chronic Immune Thrombocytopenia and Pediatric Patients 6 Years and Older with Persistent or Chronic Immune Thrombocytopenia

Platelet Count	Dose Adjustment or Action
Less than $50 \times 10^9/L$ after at least 2 weeks of DOPTELET tablets	<ul style="list-style-type: none"> • Increase <i>One Dose Level</i> per Table 3. • Wait 2 weeks to assess the effects of this regimen and any subsequent dose adjustments.
Between $200 \times 10^9/L$ and $400 \times 10^9/L$	<ul style="list-style-type: none"> • Decrease <i>One Dose Level</i> per Table 3. • Wait 2 weeks to assess the effects of this regimen and any subsequent dose adjustments.
Greater than $400 \times 10^9/L$	<ul style="list-style-type: none"> • Stop DOPTELET tablets. • Increase platelet monitoring to twice weekly. • When platelet count is less than $150 \times 10^9/L$, decrease <i>One Dose Level</i> per Table 3 and reinitiate therapy.
Less than $50 \times 10^9/L$ after 4 weeks of DOPTELET 40 mg (2 tablets) once daily	<ul style="list-style-type: none"> • Discontinue DOPTELET tablets.
Greater than $400 \times 10^9/L$ after 2 weeks of DOPTELET 20 mg (1 tablet) weekly	<ul style="list-style-type: none"> • Discontinue DOPTELET tablets.

Table 3: DOPTELET Tablet Dose Levels for Titration in Adult Patients with Chronic Immune Thrombocytopenia and Pediatric Patients 6 Years and Older with Persistent or Chronic Immune Thrombocytopenia

Dosage	Dose Level
40 mg (2 tablets) orally Once Daily	6
40 mg (2 tablets) orally Three Times a Week AND 20 mg (1 tablet) orally on the Four Remaining Days of Each Week	5
20 mg (1 tablet) orally Once Daily*	4
20 mg (1 tablet) orally Three Times a Week	3
20 mg (1 tablet) orally Twice a Week OR 40 mg Once Weekly	2
20 mg (1 tablet) orally Once Weekly	1

*Initial dosage regimen for all patients *except* those taking *Moderate or Strong Dual Inducers or Moderate or Strong Dual Inhibitors of CYP2C9 and CYP3A4*.

Discontinuation:

- Discontinue DOPTELET tablets if the platelet count does not increase to greater than or equal to $50 \times 10^9/L$ after 4 weeks of dosing at the maximum dose of 40 mg (2 tablets) once daily.

- Discontinue DOPTELET tablets if the platelet count is greater than $400 \times 10^9/L$ after 2 weeks of dosing at 20 mg (1 tablet) once weekly.

2.4 Recommended Initial Dosage of DOPTELET Tablets with Concomitant Moderate or Strong Dual Inducers or Inhibitors of CYP2C9 and CYP3A4

The recommended initial dosages of DOPTELET tablets with concomitant moderate or strong dual inducers or inhibitors of CYP2C9 and CYP3A4 in adult patients with chronic immune thrombocytopenia and pediatric patients 6 years and older with persistent or chronic immune thrombocytopenia are summarized in [Table 4](#).

Table 4: DOPTELET Tablets Recommended Initial Dosage with Concomitant Moderate or Strong Dual Inducers or Inhibitors of CYP2C9 and CYP3A4 for Adult Patients with Chronic Immune Thrombocytopenia and Pediatric Patients 6 Years and Older with Persistent or Chronic Immune Thrombocytopenia

Concomitant Medications	Recommended Initial Dosage
Moderate or strong dual inhibitors of CYP2C9 and CYP3A4	20 mg (1 tablet) orally three times a week
Moderate or strong dual inducers of CYP2C9 and CYP3A4	40 mg (2 tablets) orally once daily

2.5 Recommended Dosage of DOPTELET SPRINKLE for Patients 1 Year to Less than 6 Years with Persistent or Chronic Immune Thrombocytopenia

Use the lowest dose of DOPTELET SPRINKLE needed to achieve and maintain a platelet count greater than or equal to $50 \times 10^9/L$ as necessary to reduce the risk for bleeding. Dose adjustments are based on platelet count response. Do not use DOPTELET SPRINKLE to normalize platelet counts.

Initial Dosage:

- Begin DOPTELET SPRINKLE at an initial dosage of 10 mg (content of 1 capsule) orally once daily with food (see [Table 6](#)).
- The recommended initial dosages of DOPTELET SPRINKLE are different for patients receiving certain concomitant medications (see [Table 7](#)).

Monitoring:

After initiating therapy with DOPTELET SPRINKLE, assess platelet counts weekly until a stable platelet count greater than or equal to $50 \times 10^9/L$ has been achieved, and then obtain platelet counts monthly thereafter. Obtain platelet counts weekly for at least 4 weeks following discontinuation of DOPTELET SPRINKLE.

Dose Adjustments:

DOPTELET SPRINKLE dose adjustments (see [Table 5](#) and [Table 6](#)) are based on the platelet count response. Do not exceed a DOPTELET SPRINKLE daily dose of 20 mg (content of 2 capsules).

Table 5: DOPTELET SPRINKLE Dose Adjustments for Patients 1 Year to Less than 6 Years with Persistent or Chronic Immune Thrombocytopenia

Platelet Count	Dose Adjustment or Action
Less than $50 \times 10^9/L$ after at least 2 weeks of DOPTELET SPRINKLE	<ul style="list-style-type: none"> • Increase <i>One Dose Level</i> per Table 6. • Wait 2 weeks to assess the effects of this regimen and any subsequent dose adjustments.
Between $200 \times 10^9/L$ and $400 \times 10^9/L$	<ul style="list-style-type: none"> • Decrease <i>One Dose Level</i> per Table 6. • Wait 2 weeks to assess the effects of this regimen and any subsequent dose adjustments.
Greater than $400 \times 10^9/L$	<ul style="list-style-type: none"> • Stop DOPTELET SPRINKLE. • Increase platelet monitoring to twice weekly. • When platelet count is less than $150 \times 10^9/L$, decrease <i>One Dose Level</i> per Table 6 and reinitiate therapy.
Less than $50 \times 10^9/L$ after 4 weeks of DOPTELET SPRINKLE 20 mg (content of 2 capsules) once daily	<ul style="list-style-type: none"> • Discontinue DOPTELET SPRINKLE.
Greater than $400 \times 10^9/L$ after 2 weeks of DOPTELET SPRINKLE 10 mg (content of 1 capsule) weekly	<ul style="list-style-type: none"> • Discontinue DOPTELET SPRINKLE.

Table 6: DOPTELET SPRINKLE Dose Levels for Titration in Pediatric Patients 1 Year to Less than 6 Years with Persistent or Chronic Immune Thrombocytopenia

Dosage	Dose Level
20 mg (content of 2 capsules) orally Once Daily	6
20 mg (content of 2 capsules) orally Three Times a Week AND 10 mg (content of 1 capsule) orally on the Four Remaining Days of Each Week	5
10 mg (content of 1 capsule) orally Once Daily*	4
10 mg (content of 1 capsule) orally Three Times a Week	3
10 mg (content of 1 capsule) orally Twice a Week OR 20 mg (content of 2 capsules) orally Once Weekly	2
10 mg (content of 1 capsule) orally Once Weekly	1

*Initial dose regimen for all patients *except* those taking *Moderate or Strong Dual Inducers* or *Moderate or Strong Dual Inhibitors* of CYP2C9 and CYP3A4.

Discontinuation:

- Discontinue DOPTELET SPRINKLE if the platelet count does not increase to greater than or equal to $50 \times 10^9/L$ after 4 weeks of dosing at the maximum dose of 20 mg (content of 2 capsules) once daily.
- Discontinue DOPTELET SPRINKLE if the platelet count is greater than $400 \times 10^9/L$ after 2 weeks of dosing at 10 mg (content of 1 capsule) once weekly.

2.6 Recommended Initial Dosage of DOPTELET SPRINKLE with Concomitant Moderate or Strong Dual Inducers or Inhibitors of CYP2C9 and CYP3A4

The recommended initial dosages of DOPTELET SPRINKLE with concomitant moderate or strong dual inducers or inhibitors of CYP2C9 and CYP3A4 in pediatric patients 1 year to less than 6 years with persistent or chronic immune thrombocytopenia are summarized in [Table 7](#).

Table 7: DOPTELET SPRINKLE Recommended Initial Dosage with Concomitant Moderate or Strong Dual Inducers or Inhibitors of CYP2C9 and CYP3A4 for Pediatric Patients 1 Year to Less than 6 Years with Persistent or Chronic Immune Thrombocytopenia

Concomitant Medications	Recommended Initial Dosage
Moderate or strong dual inhibitors of CYP2C9 and CYP3A4	10 mg (content of 1 capsule) orally three times a week
Moderate or strong dual inducers of CYP2C9 and CYP3A4	20 mg (content of 2 capsules) orally once daily

2.7 Important Preparation and Administration Instructions for DOPTELET SPRINKLE

- Open the capsules and sprinkle the granules onto a small amount of a soft food or liquid in a spoon or cup.
- Do **not** swallow the capsules whole. Do not chew or crush the granules.
- Use the entire contents of the capsules to achieve the dose.
 - The following soft foods and liquids are suitable:
 - Soft foods: applesauce; strawberry jelly; yogurt (plain)
 - Liquids: milk (whole or skim); orange juice; pediatric electrolyte solution (unflavored); water
- Mix the granules into the soft food or liquid; the granules will not dissolve.
- Consume the mixture immediately after preparation; it should not be saved for future use.
- Rinse the spoon or cup with the soft food or liquid to ensure that the full dose is administered.

2.8 Missed Dose

In the case of a missed dose of DOPTELET tablets or DOPTELET SPRINKLE, patients should take the missed dose as soon as they remember. Patients should not take two doses at one time to make up for a missed dose and should take the next dose at the usual time the next day.

3 DOSAGE FORMS AND STRENGTHS

Tablets

DOPTELET Tablets: 20 mg as round, biconvex, yellow, film-coated tablets debossed with “AVA” on one side and “20” on the other side.

Oral Granules

DOPTELET SPRINKLE: 10 mg as Size 0 capsules containing white to off-white granules. The capsules have a white opaque body and light blue opaque cap with a black arrow symbol imprinted on the body and cap.

4 CONTRAINDICATIONS

None.

5 WARNINGS AND PRECAUTIONS

5.1 Thrombotic/Thromboembolic Complications

DOPTELET is a thrombopoietin (TPO) receptor agonist and TPO receptor agonists have been associated with thrombotic and thromboembolic complications in patients with chronic liver disease or immune thrombocytopenia. In patients with chronic liver disease, thromboembolic events (portal vein thrombosis) occurred in 0.4% (1/274) of patients receiving DOPTELET. In adult patients with chronic immune thrombocytopenia, thromboembolic events (arterial or venous) occurred in 7% (9/128) of patients receiving DOPTELET.

Consider the potential increased thrombotic risk when administering DOPTELET to patients with known risk factors for thromboembolism, including genetic prothrombotic conditions (e.g., Factor V Leiden, Prothrombin 20210A, Antithrombin deficiency or Protein C or S deficiency) and acquired risk factors (e.g., antiphospholipid syndrome).

DOPTELET should not be administered to patients with chronic liver disease or immune thrombocytopenia in an attempt to normalize platelet counts. Monitor platelet counts and follow the dosing guidelines to achieve target platelet counts [see *Dosage and Administration* (2.3 and 2.5)]. Monitor patients receiving DOPTELET for signs and symptoms of thromboembolic events and institute treatment promptly.

6 ADVERSE REACTIONS

The following clinically significant adverse reactions are discussed in detail in other sections of the labeling:

- Thrombotic/Thromboembolic Complications [see *Warnings and Precautions* (5.1)]

6.1 Clinical Trial Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Patients with Chronic Liver Disease

The safety of DOPTELET was evaluated in two international, identically designed, randomized, double-blind, placebo-controlled trials, ADAPT-1 and ADAPT-2, in which 430 patients with chronic liver disease and thrombocytopenia received either DOPTELET (n=274) or placebo (n=156) daily for 5 days prior to a scheduled procedure, and had 1 post-dose safety assessment. Patients were divided into two groups based on their mean platelet count at baseline:

- Low Baseline Platelet Count Cohort (less than $40 \times 10^9/L$) who received DOPTelet 60 mg once daily for 5 days
- High Baseline Platelet Count Cohort (40 to less than $50 \times 10^9/L$) who received DOPTelet 40 mg once daily for 5 days

The majority of patients were males (65%) and median subject age was 58 years (ranging from 19-86 years of age). The racial and ethnic distribution was White (60%), Asian (33%), Black (3%) and Other (3%).

The most common adverse reactions (those occurring in $\geq 3\%$ of patients) in the DOPTelet-treated groups (60 mg or 40 mg) across the pooled data from the two trials are summarized in [Table 8](#).

Table 8: Adverse Reactions with a Frequency $\geq 3\%$ in Patients with Chronic Liver Disease Treated with DOPTelet – Pooled Data ADAPT-1 and ADAPT-2

Adverse Reactions	Low Baseline Platelet Count Cohort ($<40 \times 10^9/L$)		High Baseline Platelet Count Cohort (≥ 40 to $<50 \times 10^9/L$)		Combined Baseline Platelet Count Cohorts ($<50 \times 10^9/L$)	
	DOPTelet 60 mg (N=159) %	Placebo (N=91) %	DOPTelet 40 mg (N=115) %	Placebo (N=65) %	Total DOPTelet (N=274) %	Total Placebo (N=156) %
Pyrexia	11	9	8	9	10	9
Abdominal Pain	6	7	7	6	7	6
Nausea	6	8	7	6	7	7
Headache	4	8	7	5	6	6
Fatigue	4	4	3	2	4	3
Edema Peripheral	3	2	4	2	3	2

For the Low Baseline Platelet Count Cohort, the incidence of serious adverse reactions was 7% (11/159) in the 60 mg DOPTelet treatment group. For the High Baseline Platelet Count Cohort, the incidence of serious adverse reactions was 8% (9/115) in the 40 mg DOPTelet treatment group. The most common serious adverse reaction reported with DOPTelet was hyponatremia. Two DOPTelet-treated patients (0.7%) developed hyponatremia.

Adverse reactions resulting in discontinuation of DOPTelet were anemia, pyrexia, and myalgia; each was reported in a single (0.4%) patient in the DOPTelet (60 mg) treatment group.

Adult Patients with Chronic Immune Thrombocytopenia

The safety of DOPTelet was evaluated in four clinical trials in adult patients with chronic immune thrombocytopenia: two Phase 3 trials (one randomized, double-blind, placebo-controlled trial, and one randomized, double-blind, active-controlled trial) and two Phase 2 trials (one randomized, double-blind, placebo-controlled, dose-ranging trial, and one open-label extension trial) in 161 patients with chronic immune thrombocytopenia in both the double-blind and open-label extension phases.

The pooled safety data from these four clinical trials includes 128 patients who received 2.5 to 40 mg of DOPTelet once daily for a median duration of exposure of 29.1 weeks and had 1 post-dose safety assessment.

The majority of patients were female (63%) and median subject age was 50.5 years (ranging from 18-88 years of age). The racial and ethnic distribution was White (84%), Black (6%), Asian (6%) and Other (6%).

The most common adverse reactions (those occurring in $\geq 10\%$ of patients) in the DOPTelet-treated patients across the pooled safety data from the four trials are summarized in [Table 9](#).

Table 9: Adverse Reactions with a Frequency $\geq 10\%$ in Adult Patients with Chronic Immune Thrombocytopenia Treated with DOPTelet – Pooled Data from Clinical Trials

Adverse Reactions	DOPTelet (N=128) %	Placebo (N=22) %
Headache	31	14
Fatigue	28	9
Contusion	26	18
Epistaxis	19	18
Upper Respiratory Tract Infection	15	5
Arthralgia	13	0
Gingival Bleeding	13	0
Petechiae	11	9
Nasopharyngitis	10	0

The incidence of serious adverse reactions was 9% (12/128) in the DOPTelet treatment group. Serious adverse reactions reported in more than 1 individual DOPTelet-treated patient included headache, occurring in 1.6% (2/128).

Adverse reactions resulting in discontinuation of DOPTelet that were reported in more than 1 patient included headache, occurring in 1.6% (2/128).

Pediatric Patients with Persistent or Chronic Immune Thrombocytopenia

The data described below reflect median exposure to DOPTelet of 12 weeks for 54 pediatric patients (≥ 1 to < 18 years of age) with persistent or chronic immune thrombocytopenia across the core phase of one double-blind, placebo-controlled trial [see *Clinical Studies (14.3)*].

[Table 10](#) presents the most common adverse reactions (experienced by greater than or equal to 10% of pediatric patients 1 year and older receiving DOPTelet) with a higher incidence for DOPTelet versus placebo.

Table 10: Adverse Reactions in Pediatric Patients with Persistent or Chronic Immune Thrombocytopenia Treated with DOPTELET^a

Adverse Reactions	DOPTELET (N=54) %	Placebo (N=21) %
Viral Infection ^b	20	5
Nasopharyngitis	20	10
Cough	17	0
Pyrexia	17	0
Oropharyngeal Pain	13	0

^a Adverse reactions that occurred in $\geq 10\%$ of DOPTELET-treated patients and $\geq 2\%$ more than placebo-treated patients.

^b Viral infection includes viral upper respiratory infection, viral infection, COVID-19, parainfluenza virus infection, and rhinovirus infection.

Two patients experienced serious adverse reactions: thrombocytosis and headache. Two patients experienced adverse reactions resulting in discontinuation of DOPTELET: vomiting and headache (in one patient) and leukocytosis (in one patient).

6.2 Postmarketing Experience

The following adverse reactions have been identified during post approval use of DOPTELET. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Immune System Disorders: Hypersensitivity reactions including pruritus, rash, choking sensation, erythema, pharyngeal edema, pruritus generalized, rash macular, swelling face, and swollen tongue.

7 DRUG INTERACTIONS

7.1 Effect of Other Drugs on DOPTELET in Patients with Persistent or Chronic Immune Thrombocytopenia

Moderate or Strong Dual Inhibitors of CYP2C9 and CYP3A4

Concomitant use with a moderate or strong dual inhibitor of CYP2C9 and CYP3A4 increases avatrombopag AUC [see *Clinical Pharmacology (12.3)*], which may increase the risk of DOPTELET toxicities. Reduce the starting dosage of DOPTELET when used concomitantly with a moderate or strong dual inhibitor of CYP2C9 and CYP3A4 (see [Table 4](#) and [Table 7](#)) [see *Dosage and Administration (2.4 and 2.6)*].

In patients starting moderate or strong dual inhibitors of CYP2C9 and CYP3A4 while receiving DOPTELET, monitor platelet counts and adjust DOPTELET dose as necessary (see [Table 2](#) and [Table 3](#); and [Table 5](#) and [Table 6](#)) [see *Dosage and Administration (2.3 and 2.5)*].

Moderate or Strong Dual Inducers of CYP2C9 and CYP3A4

Concomitant use with a moderate or strong dual inducer of CYP2C9 and CYP3A4 decreases avatrombopag AUC [see *Clinical Pharmacology (12.3)*], which may reduce DOPTELET efficacy. Increase the recommended starting dosage of DOPTELET when used concomitantly with a moderate or strong dual inducer of CYP2C9 and CYP3A4 (see [Table 4](#) and [Table 7](#)) [see *Dosage and Administration (2.4 and 2.6)*].

In patients starting moderate or strong dual inducers of CYP2C9 and CYP3A4 while receiving DOPTELET, monitor platelet counts and adjust DOPTELET dose as necessary (see [Table 2](#) and [Table 3](#); and [Table 5](#) and [Table 6](#)) [see *Dosage and Administration (2.3 and 2.5)*].

Patients with Chronic Liver Disease

No dosage adjustments are required for patients with chronic liver disease.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Based on findings from animal reproduction studies, DOPTELET may cause fetal harm when administered to a pregnant woman (see [Data](#)). The available data on DOPTELET in pregnant women are insufficient to inform a drug-associated risk of adverse developmental outcomes. In animal reproduction studies, oral administration of avatrombopag resulted in adverse developmental outcomes when administered during organogenesis in rabbits and during organogenesis and the lactation period in rats. However, these findings were observed at exposures based on an AUC substantially higher than the AUC observed in patients at the maximum recommended dose of 60 mg once daily. Advise pregnant women of the potential risk to a fetus.

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and of miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively.

Data

Animal Data

In embryo-fetal development studies, avatrombopag was administered during organogenesis at doses of 100, 300, and 1000 mg/kg/day in rats and doses of 100, 300, and 600 mg/kg/day in rabbits. Minimal decreases in fetal weights were observed in rats at the maternally toxic dose of 1000 mg/kg/day with exposures 190 times the human exposure based on AUC. Spontaneous abortions were observed at all doses tested in rabbits and were associated with decreased body weights and food consumption at 300 and 600 mg/kg/day; exposures at the lowest dose of 100 mg/kg/day were 10 times the AUC in patients at the maximum recommended dose of 60 mg once daily. There were no embryo-fetal effects in rats administered avatrombopag at doses up to 100 mg/kg/day (53 times the human exposure based on AUC) or rabbits administered avatrombopag at doses up to 600 mg/kg (35 times the human exposure based on AUC).

In pre- and postnatal development studies in rats, avatrombopag was administered during both the organogenesis and lactation periods at doses ranging from 5 to 600 mg/kg/day. Doses of 100, 300, and 600 mg/kg/day caused maternal toxicity leading to total litter losses, decreased body weight in pups, and increased pup mortality, with the majority of the pup mortality occurring from postnatal days 14 to 21. At a dose of 50 mg/kg/day that did not produce clear maternal toxicity, avatrombopag caused increased pup mortality from postnatal days 4 to 21, and mortality continued through postnatal day 25. The 50 mg/kg/day dose also decreased body weight gain in the pups, resulting in a delay in sexual maturation. There were no effects on

behavioral or reproductive functions in the offspring. The 50 mg/kg/day dose resulted in maternal exposures 43 times and pup exposures approximately 3 times the AUC observed in patients at the maximum recommended dose of 60 mg once daily.

8.2 Lactation

Risk Summary

There is no information regarding the presence of avatrombopag in human milk, the effects on the breastfed child, or the effects on milk production. Avatrombopag was present in the milk of lactating rats. When a drug is present in animal milk, it is likely the drug will be present in human milk. Due to the potential for serious adverse reactions in a breastfed child from DOPTOLET, breastfeeding is not recommended during treatment with DOPTOLET and for at least 2 weeks after the last dose (*see Clinical Considerations*).

Clinical Considerations

Minimizing Exposure

A lactating woman receiving DOPTOLET for brief periods, such as prior to an invasive procedure, should interrupt breastfeeding and pump and discard breastmilk during treatment and for two weeks after the last dose of DOPTOLET in order to minimize exposure to a breastfed child. Advise lactating women receiving chronic DOPTOLET therapy not to breastfeed during treatment with DOPTOLET and for at least 2 weeks after the last dose.

8.4 Pediatric Use

The safety and effectiveness of DOPTOLET tablet for the treatment of persistent or chronic ITP have been established in pediatric patients aged 6 years and older. The safety and effectiveness of DOPTOLET SPRINKLE for the treatment of persistent or chronic ITP have been established in pediatric patients aged 1 to <6 years. Use of DOPTOLET tablet and DOPTOLET SPRINKLE for their respective populations is supported by evidence from an adequate and well-controlled study in pediatric patients aged 1 year and older [*see Adverse Reactions (6.1), Clinical Pharmacology (12.3), and Clinical Studies (14.3)*].

Juvenile Animal Toxicity Data

In a 10-week juvenile toxicology study in rats, avatrombopag was administered at doses ranging from 20 to 300 mg/kg/day. There was no test article-related mortality and there were no clinical signs at doses up to 300 mg/kg/day. In the stomach, dose-dependent degeneration, regenerative hyperplasia, and atrophy of the glandular epithelium occurred at 100 and 300 mg/kg/day; exposures at 100 mg/kg/day in male rats were 14 times the AUC in patients at the highest recommended dose of 60 mg once daily. An increased incidence of background focal mineralization was also observed in the kidneys of females at 300 mg/kg/day (female rat exposure was 50 times the human exposure based on AUC at the 60 mg daily dose).

8.5 Geriatric Use

There were 112 patients 65 years of age and older in the clinical studies for Chronic Liver Disease and Chronic Immune Thrombocytopenia [*see Clinical Studies (14.1 and 14.2)*]. Of the total number of DOPTOLET-treated patients in these studies, 71 (23%) were 65 years of age and older, while 12 (4%) were 75 years of age and older. Clinical studies of DOPTOLET did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients.

10 OVERDOSAGE

In the event of overdose, platelet count may increase excessively and result in thrombotic or thromboembolic complications. Closely monitor the patient and platelet count. Treat thrombotic complications in accordance with standard of care.

No antidote for DOPTLET overdose is known.


Hemodialysis is not expected to enhance the elimination of DOPTLET because avatrombopag is only approximately 6% renally excreted and is highly bound to plasma proteins.

Consider contacting the Poison Help line (1-800-222-1222) or a medical toxicologist for additional overdose management recommendations.

11 DESCRIPTION

The active ingredient in DOPTLET is avatrombopag maleate, a thrombopoietin receptor agonist. The chemical name of avatrombopag maleate is 4-piperidinecarboxylic acid, 1-[3-chloro-5-[[[4-(4-chloro-2-thienyl)-5-(4-cyclohexyl-1-piperazinyl)-2-thiazolyl]amino]carbonyl]-2-pyridinyl]-, (2Z)-2-butenedioate (1:1). It has the molecular formula $C_{29}H_{34}Cl_2N_6O_3S_2 \cdot C_4H_4O_4$. The molecular weight is 765.73.

The structural formula is:



The aqueous solubility of avatrombopag maleate at various pH levels indicates that the drug substance is practically insoluble at pH 1 to 11.

DOPTLET is provided as an immediate-release tablet and as a sprinkle capsule that contains oral granules.

Each DOPTLET tablet contains 20 mg avatrombopag (equivalent to 23.6 mg of avatrombopag maleate) and the following inactive ingredients: colloidal silicon dioxide, crospovidone, lactose monohydrate, magnesium stearate, and microcrystalline cellulose. Coating film: ferric oxide yellow, polyethylene glycol, polyvinyl alcohol, talc, and titanium dioxide.

Each DOPTLET SPRINKLE capsule contains 10 mg avatrombopag (equivalent to 11.8 mg of avatrombopag maleate) and the following inactive ingredients: crospovidone Type A, magnesium stearate, mannitol, microcrystalline cellulose, and sodium lauryl sulfate. Capsule shells: Hypromellose.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Avatrombopag is an orally bioavailable, small molecule TPO receptor agonist that stimulates proliferation and differentiation of megakaryocytes from bone marrow progenitor cells, resulting in an increased production of platelets. Avatrombopag does not compete with TPO for binding to the TPO receptor and has an additive effect with TPO on platelet production.

12.2 Pharmacodynamics

Platelet Response

DOPTELET tablet administered to adult patients resulted in dose- and exposure-dependent elevations in platelet counts. The onset of the platelet count increase was observed within 3 to 5 days of the start of treatment, with peak effect after 10 to 13 days. Post treatment, platelet counts decreased gradually, returning to near baseline values.

Cardiac Electrophysiology

At exposures similar to that achieved at the 40 mg and 60 mg tablet dose, DOPTELET did not prolong the QT interval to any clinically relevant extent. Mean QTc prolongation effects >20 ms are not anticipated with the highest recommended therapeutic dosing regimen based on analysis of data from the pooled clinical trials in patients with chronic liver disease.

12.3 Pharmacokinetics

The pharmacokinetics of avatrombopag were determined following administration of DOPTELET tablets, and pharmacokinetic parameters are presented as geometric mean (CV%) unless otherwise specified. Avatrombopag demonstrated dose-proportional pharmacokinetics after single doses from 10 mg (0.5 times the lowest approved dosage) to 80 mg (1.3 times the highest recommended dosage). Healthy subjects administered 40 mg of avatrombopag had a geometric mean (%CV) maximal concentration (C_{max}) of 166 (84%) ng/mL and area under the time-concentration curve extrapolated to infinity (AUC_{0-inf}) of 4198 (83%) ng.hr/mL. The pharmacokinetics of avatrombopag were similar in both healthy subjects and the chronic liver disease population.

The relative bioavailability of DOPTELET SPRINKLE compared to DOPTELET tablet has not been studied. However, a prototype of DOPTELET SPRINKLE demonstrated a 22% higher geometric mean C_{max} and a 38% higher geometric mean AUC for avatrombopag compared to DOPTELET tablet, both administered to healthy adult subjects under the fed condition.

[Table 11](#) summarizes avatrombopag exposure in a virtual adult population and in pediatric patients with persistent or chronic immune thrombocytopenia.

Table 11: Summary of Avatrombopag Exposure in a Virtual Adult Population and Pediatric Patients with Persistent or Chronic Immune Thrombocytopenia

Cohort	Adults 20 mg daily DOPTELET tablet	≥12 to <18 years N=21 20 mg daily DOPTELET tablet	≥6 to <12 years N=20 20 mg daily DOPTELET tablet	≥1 to <6 years N=12 10 mg daily DOPTELET SPRINKLE
C_{max,ss} (ng/mL)				
Geometric mean (CV)	165 (61.3)	168 (36.7)	267 (25.7)	221 (35.6)
AUC_{ss} (ng × h/mL)				
Geometric mean (CV)	3277 (62.2)	3372 (41.5)	4942 (28.7)	4030 (38.6)

AUC_{ss} = area under the concentration time curve at steady-state; C_{max,ss} = maximal concentration at steady-state; CV = coefficient of variation expressed as a percent; N = number of individuals.

Absorption

The median time to maximal concentration (T_{max}) occurred at 5 to 6 hours post-dose.

Effect of Food

Avatrombopag AUC_{0-inf} and C_{max} were not affected when DOPTELET tablets were co-administered with a low-fat meal (500 calories, 3 g fat, 15 g protein, and 108 g carbohydrates) or a high-fat meal (918 calories, 59 g fat, 39 g protein, and 59 g carbohydrates). The variability of avatrombopag exposure was reduced by 40% to 60% with food. The T_{max} of avatrombopag was delayed by 0 to 2 hours when DOPTELET tablets were administered with a low-fat or high-fat meal (median T_{max} range 5 to 8 hours) compared to the fasted state.

Geometric mean exposure increased by approximately 19% and 32% for C_{max} and AUC, respectively, when a prototype of DOPTELET SPRINKLE was administered in the fed state compared to the fasted state, and between-subject variability estimates for AUC and C_{max} were decreased after administration in the fed state.

Distribution

Avatrombopag has an estimated mean apparent volume of distribution (%CV) of 180 L (25%). Avatrombopag is greater than 96% bound to human plasma proteins.

Elimination

The mean plasma elimination half-life (%CV) of avatrombopag is approximately 19 hours (19%). The mean (%CV) of the apparent clearance of avatrombopag is estimated to be 6.9 L/hr (29%).

Metabolism

Avatrombopag is primarily metabolized by cytochrome P450 CYP2C9 and CYP3A4.

Excretion

Fecal excretion accounted for 88% of the administered dose, with 34% of the dose excreted as unchanged avatrombopag. Only 6% of the administered dose was found in urine.

Specific Populations

Age (1-86 years), sex, race [Whites, African-Americans, and East Asians (i.e., Japanese, Chinese and Koreans)], and any hepatic impairment (Child-Turcotte-Pugh (CTP) grade A, B, and C, or Model for End-Stage Liver Disease (MELD) score 4-23) and mild to moderate renal impairment ($CL_{cr} \geq 30$ mL/min) did not have clinically meaningful effects on the pharmacokinetics of avatrombopag.

Lower body weight was associated with higher avatrombopag C_{max} and AUC following the proposed starting dose in patients aged 6 years and older weighing 16.3 to 175 kg. Pediatric patients aged 1 to less than 6 years had 34% higher C_{max} and 23% higher AUC following administration of 10 mg daily DOPTelet SPRINKLE compared to adults who received 20 mg daily DOPTelet tablets.

The effect of severe renal impairment ($CL_{cr} < 30$ mL/min, Cockcroft-Gault), including patients requiring hemodialysis, on avatrombopag pharmacokinetics is unknown.

Drug Interactions

Clinical Studies

Table 12 summarizes the effect of other drugs on the pharmacokinetics of avatrombopag.

Table 12: Drug Interactions: Changes in Pharmacokinetics of Avatrombopag in the Presence of Co-Administered Drug

Co-administered Drug*	Geometric Mean Ratio (90% CI) of Avatrombopag PK with/without Co-administered Drug [No Effect=1.00]	
	AUC _{0-inf}	C _{max}
Strong CYP3A Inhibitor		
Itraconazole	1.37 (1.10, 1.72)	1.07 (0.86, 1.35)
Moderate CYP3A and CYP2C9 Inhibitor		
Fluconazole	2.16 (1.71, 2.72)	1.17 (0.96, 1.42)
Moderate CYP2C9 and Strong CYP3A Inducer		
Rifampin	0.57 (0.47, 0.62)	1.04 (0.88, 1.23)
P-gp Inhibitor		
Cyclosporine	0.83 (0.65, 1.04)	0.66 (0.54, 0.82)
P-gp and Moderate CYP3A Inhibitor		
Verapamil	1.61 (1.21, 2.15)	1.26 (0.96, 1.66)

*at steady state, except for cyclosporine which was administered as a single dose.

In Vitro Studies Where Drug Interaction Potential Was Not Further Evaluated Clinically

CYP enzymes: Avatrombopag does not *inhibit* CYP1A, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6, CYP2E1 or CYP3A, does not *induce* CYP1A, CYP2B6, CYP2C or CYP3A, and *weakly induces* CYP2C8 and CYP2C9.

Transporter systems: Avatrombopag inhibits organic anion transporter (OAT) 3 and breast cancer resistance protein (BCRP), but not organic anion transporter polypeptide (OATP) 1B1 or 1B3, organic cation transporter (OCT) 2 or OAT1.

Avatrombopag is not a substrate for OATP1B1, OATP1B3, OCT2, OAT1 or OAT3.

12.5 Pharmacogenomics

The CYP2C9*2 and CYP2C9*3 loss-of-function polymorphisms result in reduced CYP2C9 enzymatic activity. In a pooled pharmacogenomic analysis of avatrombopag studies, subjects heterozygous for CYP2C9 loss-of-function polymorphisms (intermediate metabolizers [n=24]) had approximately 1.4-fold higher exposure and subjects homozygous for CYP2C9 loss-of-function polymorphisms (poor metabolizers [n=2]) had approximately 2-fold higher exposure compared to subjects wild-type for CYP2C9 (normal metabolizers [n=94]).

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

In two-year carcinogenicity studies, avatrombopag was administered orally at doses of 20, 60, and 160 mg/kg/day in mice and doses of 20, 50, and 160 mg/kg/day in rats. Avatrombopag induced a statistically significant increase in neuroendocrine cell (enterochromaffin-like cell, ECL cell) gastric tumors (carcinoids) in the stomach at 160 mg/kg in female rats. The 160 mg/kg/day dose resulted in exposures 117 times the AUC observed in patients at the maximum recommended dose of 60 mg once daily. The gastric carcinoids were considered likely due to prolonged hypergastrinemia observed in toxicity studies. Hypergastrinemia-related gastric carcinoids in rodents are generally considered to be of low risk or relevance to humans.

Avatrombopag was not mutagenic in an in vitro bacterial reverse mutation (Ames) assay or clastogenic in an in vitro human lymphocyte chromosomal aberrations assay or in an in vivo rat bone marrow micronucleus assay.

Avatrombopag did not affect fertility or early embryonic development in male rats at exposures 22 times, or in female rats at exposures 114 times, the AUC observed in patients at the maximum recommended dose of 60 mg once daily.

14 CLINICAL STUDIES

14.1 Patients with Chronic Liver Disease

The efficacy of DOPTelet for the treatment of thrombocytopenia in patients with chronic liver disease who are scheduled to undergo a procedure was established in 2 identically-designed multicenter, randomized, double-blind, placebo-controlled trials (ADAPT-1 [NCT01972529] and ADAPT-2 [NCT01976104]). In each trial, patients were assigned to the Low Baseline Platelet Count Cohort ($<40 \times 10^9/L$) or the High Baseline Platelet Count Cohort (≥ 40 to $<50 \times 10^9/L$) based on their platelet count at baseline. Patients were then randomized in a 2:1 ratio to either DOPTelet or placebo. Patients were stratified according to hepatocellular cancer (HCC) status and risk of bleeding associated with the elective procedure (low, moderate, or high). Patients undergoing neurosurgical interventions, thoracotomy, laparotomy or organ resection were not eligible for enrollment.

Patients in the Low Baseline Platelet Count Cohort received 60 mg DOPTelet or matching placebo once daily for 5 days, and patients in the High Baseline Platelet Count Cohort received 40 mg DOPTelet or matching

placebo once daily for 5 days. Eligible patients were scheduled to undergo their procedure (low, moderate, or high bleeding risk) 5 to 8 days after their last dose of treatment. Patient populations were similar between the pooled Low and High Baseline Platelet Count Cohorts and consisted of 66% male and 35% female; median age 58 years and 61% White, 34% Asian, and 3% Black.

In ADAPT-1, a total of 231 patients were randomized, 149 patients were treated with DOPTelet and 82 patients were treated with placebo. In the Low Baseline Platelet Count Cohort, the mean baseline platelet count for the DOPTelet-treated group was $31.1 \times 10^9/L$ and for the placebo-treated patients was $30.7 \times 10^9/L$. In the High Baseline Platelet Count Cohort, the mean baseline platelet count for the DOPTelet-treated patients was $44.3 \times 10^9/L$ and for placebo-treated patients was $44.9 \times 10^9/L$.

In ADAPT-2, a total of 204 patients were randomized, 128 patients were treated with DOPTelet, and 76 patients were treated with placebo. In the Low Baseline Platelet Count Cohort, the mean baseline platelet count for the DOPTelet-treated group was $32.7 \times 10^9/L$ and for the placebo-treated patients was $32.5 \times 10^9/L$. In the High Baseline Platelet Count Cohort, the mean baseline platelet count for the DOPTelet-treated patients was $44.3 \times 10^9/L$ and for the placebo-treated patients was $44.5 \times 10^9/L$.

Across both baseline platelet count cohorts and the avatrombopag and placebo treatment groups, patients underwent a broad spectrum of types of scheduled procedures that ranged from low to high bleeding risk. Overall, the majority of patients (60.8% [248/430] subjects) in all treatment groups underwent low bleeding risk procedures, 17.2% (70/430) of patients underwent procedures associated with moderate bleeding risk, and 22.1% (90/430) of subjects underwent procedures associated with high bleeding risk. The proportions of patients undergoing low, moderate, and high-risk procedures were similar between the avatrombopag and placebo treatment groups.

The major efficacy outcome was the proportion of patients who did not require a platelet transfusion or any rescue procedure for bleeding after randomization and up to 7 days following an elective procedure. Additional secondary efficacy outcomes were the proportion of patients who achieved platelet counts of $\geq 50 \times 10^9/L$ on the day of procedure, and the change in platelet count from baseline to procedure day.

Responders were defined as patients who did not require a platelet transfusion or any rescue procedure for bleeding after randomization and up to 7 days following a scheduled procedure. The following were considered rescue therapies to manage the risk of bleeding associated with a procedure: whole blood transfusion, packed red blood cell (RBC) transfusion, platelet transfusion, fresh frozen plasma (FFP) or cryoprecipitate administration, Vitamin K, desmopressin, recombinant activated factor VII, aminocaproic acid, tranexamic acid, or surgical or interventional radiology procedures performed to achieve hemostasis and control blood loss. In both baseline platelet count cohorts, patients in the DOPTelet treatment groups had a greater proportion of responders than the corresponding placebo treatment groups that was both clinically meaningful and statistically significant as detailed in [Table 13](#).

Table 13: Proportion of Patients Not Requiring a Platelet Transfusion or Any Rescue Procedure for Bleeding by Baseline Platelet Count Cohort and Treatment Group – ADAPT-1 and ADAPT-2

Low Baseline Platelet Count Cohort ($<40 \times 10^9/L$)				
Category	ADAPT-1		ADAPT-2	
	DOPTELET 60 mg (n=90)	Placebo (n=48)	DOPTELET 60 mg (n=70)	Placebo (n=43)
Responders <i>95% CI</i> ^a	66% (56, 75)	23% (11, 35)	69% (58, 79)	35% (21, 49)
Difference of Proportion vs. Placebo ^b <i>95% CI</i> ^c	43% (27, 58)		34% (16, 52)	
p-value ^d	<0.0001		0.0006	
High Baseline Platelet Count Cohort (≥ 40 to $<50 \times 10^9/L$)				
Category	ADAPT-1		ADAPT-2	
	DOPTELET 40 mg (n=59)	Placebo (n=34)	DOPTELET 40 mg (n=58)	Placebo (n=33)
Responders <i>95% CI</i> ^a	88% (80, 96)	38% (22, 55)	88% (80, 96)	33% (17, 49)
Difference of Proportion vs. Placebo ^b <i>95% CI</i> ^c	50% (32, 68)		55% (37, 73)	
p-value ^d	<0.0001		<0.0001	

a. Two-sided 95% confidence interval based on normal approximation.

b. Difference of Proportion vs. placebo = Proportion of Responders for DOPTELET – Proportion of Responders for placebo.

c. 95% confidence interval calculated based on normal approximation.

d. By Cochran-Mantel-Haenszel Testing stratified by bleeding risk for the procedure.

In addition, both trials demonstrated a higher proportion of patients who achieved the target platelet count of $\geq 50 \times 10^9/L$ on the day of procedure, a secondary efficacy endpoint, in both DOPTELET-treated groups versus the placebo-treated groups for both cohorts (Low Baseline Platelet Count Cohort – ADAPT-1: 69% vs 4%, respectively; $p < 0.0001$, ADAPT-2: 67% vs 7%, respectively; $p < 0.0001$; High Baseline Platelet Count Cohort – ADAPT-1: 88% vs 21%, respectively; $p < 0.0001$; ADAPT-2: 93% vs 39%, respectively; $p < 0.0001$). Further, both trials demonstrated a greater mean change in platelet counts from baseline to the day of the procedure, a secondary efficacy endpoint, in both DOPTELET-treated groups versus the placebo-treated groups for both cohorts (Low Baseline Platelet Count Cohort – ADAPT-1: $32 \times 10^9/L$ vs $0.8 \times 10^9/L$, respectively; $p < 0.0001$; ADAPT-2: $31.3 \times 10^9/L$ vs $3.0 \times 10^9/L$, respectively; $p < 0.0001$; High Baseline Platelet Count Cohort – ADAPT-1: $37.1 \times 10^9/L$ vs $1.0 \times 10^9/L$, respectively; $p < 0.0001$; ADAPT-2: $44.9 \times 10^9/L$ vs $5.9 \times 10^9/L$, respectively; $p < 0.0001$).

A measured increase in platelet counts was observed in both DOPTELET treatment groups over time beginning on Day 4 post-dose, that peaked on Day 10-13, decreased 7 days post-procedure, and then returned to near baseline values by Day 35.

14.2 Adult Patients with Chronic Immune Thrombocytopenia

Randomized Phase 3 Clinical Trial

The efficacy of DOPTelet in adult patients with chronic immune thrombocytopenia was evaluated in a Phase 3, multicenter, randomized, double-blind, placebo-controlled trial (NCT01438840). Patients had previously received one or more prior chronic immune thrombocytopenia therapies and had an average of screening and baseline platelet counts $<30 \times 10^9/L$. Patients were centrally stratified by splenectomy status, baseline platelet count ($\leq 15 \times 10^9/L$ or $>15 \times 10^9/L$ to $<30 \times 10^9/L$), and use of concomitant chronic immune thrombocytopenia medication, and then randomized (2:1) to receive either DOPTelet or placebo for 6 months. Patients received a starting dose of 20 mg once daily, with doses subsequently titrated based on platelet response.

Forty-nine patients were randomized, 32 to DOPTelet and 17 to placebo, with similar mean [SD] baseline platelet counts in the 2 treatment groups ($14.1 [8.6] \times 10^9/L$ and $12.7 [7.8] \times 10^9/L$, respectively). The median age was 44 years, 63% were female, and 94% were Caucasian, 4% Asian and 2% Black. The median duration of exposure was 26 weeks for DOPTelet-treated patients and 6 weeks for placebo-treated patients. The major efficacy outcome in this trial was the cumulative number of weeks in which the platelet count was $\geq 50 \times 10^9/L$ during the 6-month treatment period in the absence of rescue therapy. DOPTelet-treated patients had a longer duration of platelet counts $\geq 50 \times 10^9/L$ in the absence of rescue therapy than those who received placebo (median 12.4 [0, 25] vs 0 [0, 2] weeks, respectively, $p < 0.0001$) (see [Table 14](#)).

Table 14: Cumulative Number of Weeks of Platelet Response – Phase 3 Trial in Adult Patients with Chronic Immune Thrombocytopenia

Primary Efficacy Analysis	DOPTelet (n=32)	Placebo (n=17)
Cumulative Number of Weeks with a Platelet Response*		
Mean (SD)	12.0 (8.75)	0.1 (0.49)
Median	12.4	0.0
Min, Max	0, 25	0, 2
p-value of Wilcoxon rank sum test	<0.0001	

Max=maximum, Min=minimum, SD=Standard deviation.

*Cumulative number of weeks of platelet response is defined as the total numbers of weeks in which the platelet count was $\geq 50 \times 10^9/L$ during 6 months of treatment in the absence of rescue therapy.

In addition, a larger proportion of patients in the DOPTelet treatment group had platelet counts $\geq 50 \times 10^9/L$ at Day 8 compared to placebo (21/32; 66% vs 0/17; 0.0%, respectively; $p < 0.0001$).

14.3 Pediatric Patients with Persistent or Chronic Immune Thrombocytopenia

The efficacy and safety of DOPTelet was evaluated in pediatric patients ≥ 1 to <18 years of age with persistent or chronic immune thrombocytopenia in a randomized, double-blind, placebo-controlled trial (NCT 04516967), which included a 12-week randomized treatment phase (Core Phase).

Patients were required to have had a diagnosis of primary ITP for ≥ 6 months and had an insufficient response to at least one previous treatment, with an average of 2 baseline platelet counts less than $30 \times 10^9/L$. Patients ($n=75$) were randomized (3:1) to receive DOPTelet ($n=54$) or placebo ($n=21$). The starting dose for patients 6 years and older was 20 mg (tablet), while the starting dose for the youngest cohort was 10 mg (oral granules mixed with a soft food or liquid). Doses could be subsequently titrated based on platelet response.

Enrollment was 52% male and 48% female. The median age of patients receiving DOPTelet was 8.5 years (range 1 to 17) while the median age of patients receiving placebo was 10.0 years (range 3 to 17). Patients identified their race as White (84%), Asian (5.3%), and Other (5.3%); 5.3% did not report race. Patients identified their ethnicity as Not Hispanic or Latino (86.7%) and Hispanic or Latino (6.7%); 2.7% did not report ethnicity and 4% had unknown ethnicity.

The median baseline platelet counts were $10.4 \times 10^9/L$ in the DOPTelet group and $11.5 \times 10^9/L$ in the placebo group. The percentage of patients who had received 3 or more prior ITP therapies was 68.5% in the avatrombopag group and 66.7% in the placebo group. Most patients in the trial received at least one other TPO receptor agonist as a prior therapy (74.1% in the DOPTelet group and 71.4% in the placebo group).

The efficacy of DOPTelet in this trial was evaluated by durable platelet response, defined as the proportion of patients achieving at least 6 out of 8 weekly platelet counts $\geq 50 \times 10^9/L$ during the last 8 weeks of the 12-week Treatment Period in the Core Phase in the absence of rescue medication (Table 15).

Efficacy was also evaluated by platelet response, defined as the proportion of subjects achieving at least 2 consecutive platelet assessments $\geq 50 \times 10^9/L$ in the Core Phase in the absence of rescue medication.

Table 15: Durable Platelet Response and Platelet Response - Phase 3 Trial in Pediatric Patients with Persistent or Chronic ITP - Full Analysis Set

Endpoint	DOPTelet (N=54)	Placebo (N=21)
Durable platelet response, n%		
Yes	15 (27.8)	0
No	39 (72.2)	21 (100.0)
Difference of proportion (avatrombopag – placebo) (95% CI)	27.8 (15.8, 39.7)	
CMH (avatrombopag vs. placebo) p-value	$p=0.0077^a$	
Platelet response, n%		
Yes	44 (81.5)	0
No	10 (18.5)	21 (100.0)
Difference of proportion (avatrombopag – placebo) (95% CI)	81.5 (71.1, 91.8)	
CMH (avatrombopag vs. placebo) p-value	$p < 0.0001^a$	

CI, Confidence interval; CMH, Cochran-Mantel-Haenszel; N, Total number of subjects; n, Number of subjects; Full Analysis Set includes all randomized subjects.

^a Denotes p-value from Fisher's Exact Test, which was used in place of CMH test due to sparse number of responders in the strata.

Note: The CMH test is adjusted for age cohort and baseline platelet counts.

DOPTELET was superior to placebo in other major efficacy outcomes that evaluated platelet counts. At Day 8, 55.6% (95% CI: 41.4%, 69.1%) of DOPTELET versus no placebo patients (95% CI: 0.0%, 16.1%) had a platelet count $\geq 50 \times 10^9/L$ in the absence of rescue therapy ($p < 0.0001$). The mean percentage of weeks that subjects had a platelet count $\geq 50 \times 10^9/L$ during the Core Phase in the absence of rescue therapy was significantly higher for DOPTELET versus placebo, 48.9% (SD: 25.22%) versus 1.2% (SD: 3.92%) ($p < 0.0001$).

The proportion of subjects who required rescue therapy during the Core Phase was significantly lower ($p = 0.0008$) in the DOPTELET group (7.4% [95% CI: 2.1%, 17.9%]) than in the placebo group (42.9% [95% CI: 21.8%, 66.0%]).

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 DOPTELET Tablets

DOPTELET 20 mg tablets are supplied as round, biconvex, yellow, film-coated tablets, and debossed with “AVA” on one side and “20” on the other side.

How Supplied	Carton NDC	Blister Card NDC
Carton of one blister card with 10 tablets	NDC 71369-020-10	NDC 71369-020-11
Carton of one blister card with 15 tablets	NDC 71369-020-15	NDC 71369-020-16
Carton of two blister cards, each with 15 tablets (30 tablets total)	NDC 71369-020-30	NDC 71369-020-16

Store DOPTELET tablets at room temperature from 20°C to 25°C (68°F to 77°F), excursions permitted from 15°C to 30°C (59°F to 86°F). Store tablets in the original package.

16.2 DOPTELET SPRINKLE

DOPTELET SPRINKLE is supplied as 10 mg capsules with a white opaque body and light blue opaque cap with a black arrow symbol imprinted on the body and cap. The capsule is filled with white to off-white granules.

How Supplied	Carton and Bottle NDC
Carton with one bottle of 30 capsules (containing oral granules)	NDC 71369-010-30

Store DOPTELET SPRINKLE capsules at room temperature from 20°C to 25°C (68°F to 77°F), excursions permitted from 15°C to 30°C (59°F to 86°F).

17 PATIENT COUNSELING INFORMATION

Advise the patient or caregiver to read the FDA-approved patient labeling: Patient Information and, for DOPTELET SPRINKLE, Instructions for Use.

Prior to treatment, patients should fully understand and be informed of the following risks and considerations for DOPTELET:

Risks

Thrombotic/Thromboembolic Complications

DOPTELET is a thrombopoietin (TPO) receptor agonist and TPO receptor agonists have been associated with thrombotic and thromboembolic complications in patients with chronic liver disease or immune thrombocytopenia. Portal vein thrombosis has been reported in patients with chronic liver disease treated with TPO receptor agonists. Various thromboembolic complications (arterial and venous) have been reported in patients treated with DOPTELET [see *Warnings and Precautions (5.1)*].

Drug Interactions

DOPTELET may be affected by other drugs and may require a dose adjustment when co-administered with other drugs; therefore, advise patients to report their use of any other prescription or nonprescription medications or dietary supplements [see *Dosage and Administration (2.4 and 2.6)*, *Drug Interactions (7)*].

Pregnancy

Advise pregnant women of the potential risk to a fetus. Advise females of reproductive potential to inform their prescriber of a known or suspected pregnancy [see *Use in Specific Populations (8.1)*].

Lactation

Advise women not to breastfeed during treatment with DOPTELET and for at least 2 weeks after the last dose [see *Use in Specific Populations (8.2)*].

Administration Instructions for DOPTELET SPRINKLE

- Inform patients and caregivers to open the capsules and mix the contents with the recommended soft foods or liquids. Administer immediately after mixing. Do not swallow the capsules whole. Do not chew or crush the granules.
- Advise patients and caregivers to read and follow the Instructions for Use for DOPTELET SPRINKLE.

DOPTELET and DOPTELET SPRINKLE are not substitutable on a milligram-to-milligram basis

- Advise patients and caregivers that DOPTELET and DOPTELET SPRINKLE are not substitutable on a milligram-to-milligram basis.
- To avoid a dosing error from using the wrong dosage form, strongly advise patients and caregivers to visually inspect the product to verify the correct dosage form each time the prescription is filled.

DOPTELET is a registered trademark of AkaRx, Inc.
Manufactured for AkaRx, Inc., Morrisville, North Carolina 27560
©2025

For more information, go to www.DOPTELET.com or call 1-855-454-3887.

PI0002 R8

PATIENT INFORMATION

DOPTELET® (dop-TEL-et)
(avatrombopag) tablets, for oral use

DOPTELET® SPRINKLE (dop-TEL-et SPRINK-el)
(avatrombopag) oral granules

What are DOPTELET and DOPTELET SPRINKLE?

- DOPTELET is a prescription medicine used to treat low blood platelet counts in:
 - adults with long-lasting (chronic) liver disease (CLD) who are scheduled to have a medical or dental procedure.
 - adults with chronic immune thrombocytopenia (ITP) when other treatments have not worked well enough.
 - children 1 year and older with persistent or chronic ITP when other treatments have not worked well enough.

DOPTELET tablets and DOPTELET SPRINKLE are different dosage forms of DOPTELET. DOPTELET SPRINKLE is a capsule that contains granules that are mixed with a soft food or liquid for dosing.

DOPTELET tablets and DOPTELET SPRINKLE are not used to make platelet counts normal.

It is not known if DOPTELET tablets or DOPTELET SPRINKLE are safe and effective in children younger than 1 year of age.

Before you take DOPTELET or DOPTELET SPRINKLE, tell your healthcare provider about all of your medical conditions, including if you:

- have ever had a blood clot.
- are pregnant or plan to become pregnant. DOPTELET may harm your unborn baby. Tell your healthcare provider if you become pregnant or think you may be pregnant during treatment with DOPTELET.
- are breastfeeding or plan to breastfeed. It is not known if DOPTELET passes into your breast milk. Do not breastfeed during your treatment with DOPTELET and for at least 2 weeks after the last dose. Talk to your healthcare provider about the best way to feed your baby during this time.

Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. DOPTELET or DOPTELET SPRINKLE may affect the way other medicines work, and other medicines may affect the way DOPTELET or DOPTELET SPRINKLE works.

How should I take DOPTELET?

- DOPTELET tablets that come in a blister package are **not** the same dosage form as DOPTELET SPRINKLE that come in a bottle and **cannot** be directly substituted for each other. Contact your or your child's pharmacist or healthcare provider if you did not receive the correct dosage form.
- Take DOPTELET exactly as your healthcare provider tells you to take it.
- Your healthcare provider will tell you how much DOPTELET to take and when to start taking it.
- Your healthcare provider may change your dose of DOPTELET depending on your blood platelet counts.
- Take DOPTELET with food.
- If you take DOPTELET to treat your low blood platelet counts due to chronic liver disease before a medical or dental procedure, your healthcare provider will check your platelet count before treatment and on the day of your scheduled procedure.
- If you take DOPTELET to treat your low blood platelet counts due to persistent or chronic immune thrombocytopenia, your healthcare provider will check your platelet count before, during and for at least 4 weeks after stopping your treatment with DOPTELET.
- If you are taking DOPTELET before a scheduled medical procedure and you miss a dose, contact your healthcare provider for further dosing instructions.
- If you are taking DOPTELET for persistent or chronic immune thrombocytopenia and you miss a dose, take it as soon as you remember. Do not take 2 doses at one time to make up for a missed dose. Take your next dose at your usual scheduled time.
- If you take too much DOPTELET, call your healthcare provider or Poison Help Line at 1-800-222-1222, or go to the nearest hospital emergency room right away.

How should I give DOPTLET SPRINKLE?

- DOPTLET SPRINKLE oral granules that come in a bottle are **not** the same dosage form as DOPTLET tablets that come in a blister card and **cannot** be directly substituted for each other. Contact your child's pharmacist or healthcare provider if you did not receive the correct dosage form.
- If your child is prescribed DOPTLET SPRINKLE: **See the Instructions for Use** on how to prepare and give a dose.
- Open the capsules and sprinkle the contents onto a small amount of a soft food or liquid. Mix or stir the granules into the soft food or liquid and take right away.
- **Do not** swallow the capsules whole or the empty capsule shells.
- Do not chew or crush the oral granules.
- Give DOPTLET SPRINKLE exactly as your child's healthcare provider tells you to give it.
- Your child's healthcare provider will tell you how much DOPTLET SPRINKLE to give and when to start taking it.
- Your child's healthcare provider may change the dose of DOPTLET SPRINKLE.
- Give your child DOPTLET SPRINKLE with food.
- If your child is taking DOPTLET SPRINKLE to treat low blood platelet counts due to persistent or chronic immune thrombocytopenia, your child's healthcare provider will check your child's platelet count before, during and for at least 4 weeks after stopping treatment with DOPTLET SPRINKLE.
- If you are giving DOPTLET SPRINKLE for your child's persistent or chronic immune thrombocytopenia and miss a dose, give it as soon as you remember. Do not give 2 doses at one time to make up for a missed dose. Give the next dose at the usual scheduled time.
- If your child takes too much DOPTLET SPRINKLE, call your child's healthcare provider or Poison Help Line at 1-800-222-1222 or go to the nearest hospital emergency room right away.

What are the possible side effects of DOPTLET or DOPTLET SPRINKLE?

DOPTLET or DOPTLET SPRINKLE may cause serious side effects, including:

Blood clots. People with chronic liver disease or persistent or chronic immune thrombocytopenia and people with certain blood clotting conditions may have an increased risk of developing blood clots. Tell your healthcare provider right away if you get signs and symptoms of a blood clot, including:

- swelling, pain, or tenderness in your legs
- shortness of breath
- chest pain
- fast heartbeat
- stomach (abdominal) pain or tenderness

The most common side effects of DOPTLET when used to treat low blood platelet counts in adults with chronic liver disease (CLD) who are scheduled to have a medical or dental procedure are:

- fever
- stomach (abdominal) pain
- nausea
- headache
- tiredness
- swelling of hands or feet

The most common side effects of DOPTLET when used to treat low blood platelet counts in adults with chronic immune thrombocytopenia (ITP) are:

- headache
- tiredness
- bruising
- nosebleed
- upper respiratory infection
- joint pain
- bleeding gums
- purple or red spots on your skin
- runny nose

The most common side effects of DOPTLET or DOPTLET SPRINKLE when used to treat low blood platelet counts in children 1 year and older with persistent or chronic immune thrombocytopenia (ITP) are:

- viral infection
- runny nose
- cough
- fever
- pain in the mouth or throat

These are not all of the possible side effects of DOPTLET or DOPTLET SPRINKLE.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store DOPTLET tablets or DOPTLET SPRINKLE?

- Store DOPTLET tablets or DOPTLET SPRINKLE at room temperature from 68°F to 77°F (20°C to 25°C).
- Store DOPTLET tablets in the original package.

Keep DOPTLET tablets, DOPTLET SPRINKLE and all medicines out of the reach of children.

General information about the safe and effective use of DOPTELET or DOPTELET SPRINKLE.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use DOPTELET or DOPTELET SPRINKLE for a condition for which it was not prescribed. Do not give DOPTELET tablets or DOPTELET SPRINKLE to other people, even if they have the same symptoms that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about DOPTELET or DOPTELET SPRINKLE that is written for health professionals.

What are the ingredients in DOPTELET tablets or DOPTELET SPRINKLE?

Active ingredient: avatrombopag

Inactive ingredients in DOPTELET tablets: colloidal silicon dioxide, crospovidone, lactose monohydrate, magnesium stearate and microcrystalline cellulose. Tablet coating film: ferric oxide yellow, polyethylene glycol, polyvinyl alcohol, talc, and titanium dioxide.

Inactive ingredients in DOPTELET SPRINKLE: crospovidone Type A, magnesium stearate, mannitol, microcrystalline cellulose, and sodium lauryl sulfate. Capsule shells: Hypromellose.

DOPTELET is a registered trademark of AkaRx, Inc.

Manufactured for AkaRx, Inc., Morrisville, North Carolina 27560

For more information, go to www.DOPTELET.com or call 1-855-454-3887.

This Patient Information has been approved by the U.S. Food and Drug Administration.

Revised: 07/2025 PI0001 R8

INSTRUCTIONS FOR USE
DOPTELET® SPRINKLE (dop-TEL-et SPRINK-el)
(avatrombopag) oral granules

Read these Instructions for Use to prepare and give a dose of DOPTELET SPRINKLE correctly.

- **Important information that you need to know before giving DOPTELET SPRINKLE:**
The contents of DOPTELET SPRINKLE should be sprinkled onto a soft food or liquid, as described below.
- DOPTELET SPRINKLE that comes in a bottle is **not** the same dosage form as DOPTELET tablets that come in a blister card and these products **cannot** be directly substituted for each other. Contact your child's pharmacist or healthcare provider if your child did not receive the correct dosage form.
- DOPTELET SPRINKLE contains granules inside of a capsule.
 - The capsule must be opened, and the granules inside must be sprinkled onto a soft food or liquid to give a dose of DOPTELET SPRINKLE.
 - Do not swallow the capsule whole or the empty capsule shells.
 - Do not chew or crush the granules.
- Use all of the granules in the capsule. **Do not** use part of a capsule to try to prepare a dose.

Steps to prepare and give a dose of DOPTELET SPRINKLE:

1. Gather supplies to prepare a dose of DOPTELET SPRINKLE: capsule(s), liquid or soft food, cup or spoon, and mixing utensil (e.g., stir stick or small spoon); then wash your hands.
2. Put a small amount of soft food or liquid in a cup or spoon. The following soft foods and liquids can be used:

Soft foods

- applesauce
- strawberry jelly
- yogurt (plain)

Liquids

- milk (whole or skim)
- orange juice
- pediatric electrolyte solution (unflavored)
- water

3. Carefully open the capsule (or 2 capsules if prescribed) by pulling it apart.
 - Sprinkle the entire contents of the capsule onto the soft food or liquid that you put in the spoon or cup. Make sure that all of the granules from the capsule have been completely emptied.
4. Mix or stir the granules into the soft food or liquid. The granules will **not** dissolve.
5. Give the mixture **right away**; **do not** save for later use.
6. Rinse the spoon or cup with additional soft food or liquid to make sure that the entire dose is given.
7. Throw the empty capsule shells away in the trash.

How should I store DOPTELET SPRINKLE?

- Store DOPTELET SPRINKLE at room temperature from 68°F to 77°F (20°C to 25°C).

Keep DOPTELET SPRINKLE and all medicines out of the reach of children.

DOPTELET is a registered trademark of AkaRx, Inc.
Manufactured for AkaRx, Inc., Morrisville, North Carolina 27560
For more information, go to www.DOPTELET.com or call 1-855-454-3887.