

bel may not be the latest approved b
information, please visit <https://www>



TO OPEN PEEL FROM
ONE CORNER

Single-Dose Prefilled Pen

NDC 0597-0375-11

R_x only

CYLTEZO[®] PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL

FOR SUBCUTANEOUS USE ONLY

Contains: One sterile single-dose prefilled pen with a fixed ½ inch length needle.

Contains no preservatives.

Needle Cap Contains Dry Natural Rubber.

Recommended Dosage: See Prescribing Information.

Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light. DO NOT FREEZE.

If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use. Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days.

Write the date removed from the refrigerator

___/___/___.

Do not use beyond expiration date.

Manufactured by:
Boehringer Ingelheim Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
US License No. 2006

BL7414E



Boehringer
Ingelheim



10011164-01

LOT 1234567

EXP MMMYYYY

Label may not be the latest approved by
For more information, please visit <https://www.fda.gov>



TO OPEN PEEL FROM
ONE CORNER

Single-Dose Prefilled Syringe

NDC 0597-0370-86

R_x only

CYLTEZO[®]
(adalimumab-adbm)
Injection

40 mg/0.8 mL

FOR SUBCUTANEOUS USE ONLY

Contains: One sterile single-dose prefilled syringe with a fixed 1/2 inch length needle.

Contains no preservatives.

Needle Cap Contains Dry Natural Rubber.

Recommended Dosage: See Prescribing Information.

Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light.

DO NOT FREEZE. If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use.

Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days.

Write the date removed from the refrigerator

___/___/___

Do not use beyond expiration date.

Manufactured by:
Boehringer Ingelheim Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
US License No. 2006



Boehringer
Ingelheim



(01)00305970370864

BL7420E

10011165-02

LOT 1234567

EXP MMMYYYY

Label may not be the latest approved by
FDA. For the latest information, please visit <https://www.fda.gov>.



TO OPEN PEEL FROM
ONE CORNER

Single-Dose Prefilled Syringe

NDC 0597-0405-06

R_x only

CYLTEZO[®]
(adalimumab-adbm)
Injection

20 mg/0.4 mL

FOR SUBCUTANEOUS USE ONLY

Contains: One sterile single-dose prefilled syringe with a fixed ½ inch length needle.

Contains no preservatives.

Needle Cap Contains Dry Natural Rubber.

Recommended Dosage: See Prescribing Information.

Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light.

DO NOT FREEZE. If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use.

Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days.

Write the date removed from the refrigerator

___/___/___.

Do not use beyond expiration date.

Manufactured by:
Boehringer Ingelheim Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
US License No. 2006

BL7976D



Boehringer
Ingelheim



10011163-02

LOT 1234567

EXP MMMYYY



TO OPEN PEEL FROM
ONE CORNER

Single-Dose Prefilled Syringe

NDC 0597-0400-37

R_x only

CYLTEZO[®]
(adalimumab-adbm)
Injection

10 mg/0.2 mL

FOR SUBCUTANEOUS USE ONLY

Contains: One sterile single-dose prefilled syringe with a fixed ½ inch length needle.

Contains no preservatives.

Needle Cap Contains Dry Natural Rubber.

Recommended Dosage: See Prescribing Information.

Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light.

DO NOT FREEZE. If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use.

Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days.

Write the date removed from the refrigerator

___/___/___.

Do not use beyond expiration date.

Manufactured by:
Boehringer Ingelheim Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
US License No. 2006

BL10034C



Boehringer
Ingelheim



(01)00305970400370

10011316-02

LOT 1234567

EXP MMMYYYY



Do not use beyond the expiration date.

NDC 0597-0375-97

CYLTEZO® PEN
(adalimumab-adbm)
Injection

Citrate Free

40 mg/0.8 mL

For Subcutaneous Use Only

2 Single-Dose Prefilled Pens

Product and trademark licensed from:
Boehringer Ingelheim International GmbH
© 2023 Boehringer Ingelheim International GmbH
ALL RIGHTS RESERVED
Manufactured by:
Boehringer Ingelheim Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
Product of the USA
US License Number 2006
CT7417E



For Subcutaneous Use Only

This Carton Contains:
• 2 dose trays (each containing 1 single-dose prefilled pen with a fixed 1/2 inch length needle)
• 2 alcohol preps
• 1 Package Insert
• 1 Medication Guide
• 1 Instructions for Use

Recommended Dosage: See Prescribing Information.
Needle Cap for Pen Contains Dry Natural Rubber.

For Subcutaneous Use Only

Each Sterile Single-Dose Prefilled Pen Contains:
Adalimumab-adbm.....40 mg
Glacial acetic acid.....0.13 mg
Polysorbate 80.....0.8 mg
Sodium acetate trihydrate.....2.4 mg
Trehalose dihydrate.....65.0 mg
Water for Injection, USP

Contains no preservatives.
Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light. **DO NOT FREEZE.** If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use. Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days.
Write the date removed from the refrigerator _____.

40 mg/0.8 mL

CYLTEZO® PEN
(adalimumab-adbm)
Injection

NDC 0597-0375-97

NOTE TO PHARMACIST:
(Place Prescription Label Here)

2 Single-Dose Prefilled Pens

40 mg/0.8 mL

For Subcutaneous Use Only

CYLTEZO® PEN
(adalimumab-adbm)
Injection

NDC 0597-0375-97

For more information about CYLTEZO, including current prescribing information and Medication Guide, go to www.cyltezo.com, scan the code, or call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.



2 Single-Dose Prefilled Pens

For Subcutaneous Use Only

40 mg/0.8 mL

CYLTEZO® PEN
(adalimumab-adbm)
Injection

NDC 0597-0375-97



GTN 00305970375975
SN 12345678901234
EXP MMYYYY
LOT 1234567



10010940-02

NDC 0597-0375-97

CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

Rx only

2 Single-Dose Prefilled Pens



ATTENTION PHARMACIST:
Each Patient is Required to Receive the Enclosed Medication Guide.

The Entire Carton is to be Dispensed as a Unit.
Needle Cap for Pen Contains Dry Natural Rubber.







CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

2 Single-Dose Prefilled Pens

QUICKTIPS for USING CYLTEZO® Pen (adalimumab-adbm) Injection
For complete instructions refer to the Instructions for Use in the carton.
Do not try to inject CYLTEZO® yourself until you have been shown the right way to give the injection.

1 Choose Injection Site and Clean	2 Inspect and Uncap	3 Prepare Injection	4 Administer Dose
 <p>Abdomen or Thighs</p>		 <p>Squeeze around injection site</p> <p>Press and hold 90°</p> <p>Tip</p>	 <p>Press button 1 time and slowly count to 10</p> <p>Plunger</p>
<ul style="list-style-type: none">• Choose an area on your upper thighs or belly, except for an area 2 inches around your belly button.• Choose a different injection site each time you inject, at least 1 inch away from the previous injection site.• Clean the injection site with an alcohol wipe.	<ul style="list-style-type: none">• Inspect the pen, medicine, and expiration date.• Pull the cap straight off the pen.• Do Not twist the cap.	<ul style="list-style-type: none">• Gently squeeze the skin around the injection site and hold firmly.• Hold the pen straight up from the injection site (at a 90-degree angle).• Press and hold the tip firmly against squeezed skin at the injection site. The injection button will unlock and be ready for use.	<ul style="list-style-type: none">• Press the injection button 1 time. The 'click' sound indicates injection start.• Keep holding the pen against the squeezed skin around the injection site while you slowly count to 10.• Do Not move the pen during injection.• You will see the plunger in the window at the bottom of the pen when the injection is complete.

For more information about CYLTEZO®, call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.



Do not use beyond the expiration date.

NDC 0597-0375-53

CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only
1 Single-Dose Prefilled Pen

Product and trademark licensed from:
Boehringer Ingelheim International GmbH
© 2023 Boehringer Ingelheim International GmbH
ALL RIGHTS RESERVED
Manufactured by:
Boehringer Ingelheim Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
Product of the USA
US License Number 2006
CT7416D

Citrate Free



This Carton Contains:
• 1 dose tray containing 1 single-dose prefilled pen with a fixed 1/2 inch length needle
• 2 alcohol preps
• 1 Package Insert
• 1 Medication Guide
• 1 Instructions for Use

Recommended Dosage: See Prescribing Information.

Needle Cap for Pen Contains Dry Natural Rubber.

Contains no preservatives.

Each Sterile Single-Dose Prefilled Pen Contains:
Adalimumab-adbm.....40 mg
Glacial acetic acid.....0.13 mg
Polysorbate 80.....0.8 mg
Sodium acetate trihydrate.....2.4 mg
Trehalose dihydrate.....65.0 mg
Water for Injection, USP

Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light. DO NOT FREEZE. If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use. Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days. Write the date removed from the refrigerator.

40 mg/0.8 mL

CYLTEZO® PEN
(adalimumab-adbm)
Injection

NDC 0597-0375-53

NOTE TO PHARMACIST:
{Place Label Here}

1 Single-Dose Prefilled Pen

For Subcutaneous Use Only

40 mg/0.8 mL

Injection
(adalimumab-adbm)

NDC 0597-0375-53



GTIN
SN
12345678901234
EXP
MMYY
1234567
LOT
00305970375531



NDC 0597-0375-53

CYLTEZO® PEN
(adalimumab-adbm)
Injection

ATTENTION PHARMACIST:
Each Patient is Required to Receive the Enclosed Medication Guide.
The Entire Carton is to be Dispensed as a Unit.
Needle Cap for Pen Contains Dry Natural Rubber.

40 mg/0.8 mL

For Subcutaneous Use Only

NOT FOR SALE

Rx only

1 Single-Dose Prefilled Pen



CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

1 Single-Dose Prefilled Pen

For more information about CYLTEZO, including current prescribing information and Medication Guide, go to www.cyltezo.com, scan the code, or call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.



CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

1 Single-Dose Prefilled Pen

NDC 0597-0375-53





10010939-01

QUICKTIPS for USING CYLTEZO® Pen (adalimumab-adbm) Injection

For complete instructions refer to the Instructions for Use in the carton.

Do not try to inject CYLTEZO® yourself until you have been shown the right way to give the injection.

For more information about CYLTEZO®, call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.

1 Choose Injection Site and Clean	2 Inspect and Uncap	3 Prepare Injection	4 Administer Dose
 <p>Abdomen or Thighs</p>		 <p>Squeeze around injection site</p> <p>Press and hold 90°</p> <p>Tip</p>	 <p>Press button 1 time and slowly count to 10</p> <p>Plunger</p>
<ul style="list-style-type: none">• Choose an area on your upper thighs or belly, except for an area 2 inches around your belly button.• Choose a different injection site each time you inject, at least 1 inch away from the previous injection site.• Clean the injection site with an alcohol wipe.	<ul style="list-style-type: none">• Inspect the pen, medicine, and expiration date.• Pull the cap straight off the pen.• Do Not twist the cap.	<ul style="list-style-type: none">• Gently squeeze the skin around the injection site and hold firmly.• Hold the pen straight up from the injection site (at a 90-degree angle).• Press and hold the tip firmly against squeezed skin at the injection site. The injection button will unlock and be ready for use.	<ul style="list-style-type: none">• Press the injection button 1 time. The 'click' sound indicates injection start.• Keep holding the pen against the squeezed skin around the injection site while you slowly count to 10.• Do Not move the pen during injection.• You will see the plunger in the window at the bottom of the pen when the injection is complete.



Do not use beyond the expiration date.

NDC 0597-0375-23

CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

Citrate Free

Product and trademark licensed from:
Boehringer Ingelheim International GmbH
© 2023 Boehringer Ingelheim International GmbH
ALL RIGHTS RESERVED
Manufactured by:
Boehringer Ingelheim Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
Product of the USA
US License Number 2006
CT7418D

4 Single-Dose Prefilled Pens



For Subcutaneous Use Only

This Carton Contains:

- 4 dose trays (each containing 1 single-dose prefilled pen with a fixed 1/2 inch length needle)
- 4 alcohol prep pads
- 1 Package Insert
- 1 Medication Guide
- 1 Instructions for Use

Recommended Dosage: See Prescribing Information.

Needle Cap for Pen Contains Dry Natural Rubber.

Starter Package for Psoriasis

Write the date removed from the refrigerator _____ / ____ / ____

Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light. **DO NOT FREEZE.** If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use.

Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days.

Contains no preservatives.

Water for Injection, USP.

Trehalose dihydrate.....65.0 mg

Sodium acetate trihydrate.....2.4 mg

Polysorbate 80.....0.8 mg

Glacial acetic acid.....0.13 mg

Adalimumab-adbm.....40 mg

Each Sterile Single-Dose Prefilled Pen Contains:

40 mg/0.8 mL

CYLTEZO® PEN
(adalimumab-adbm)
Injection



NDC 0597-0375-23

NOTE TO PHARMACIST:
(Place Prescription Label Here)

Starter Package for Psoriasis

4 Single-Dose Prefilled Pens

40 mg/0.8 mL
For Subcutaneous Use Only

CYLTEZO® PEN
(adalimumab-adbm)
Injection

NDC 0597-0375-23

For more information about CYLTEZO, including current prescribing information and Medication Guide, go to www.cyltezo.com, scan the code, or call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.



Starter Package for Psoriasis

4 Single-Dose Prefilled Pens

For Subcutaneous Use Only

40 mg/0.8 mL

CYLTEZO® PEN
(adalimumab-adbm)
Injection

NDC 0597-0375-23



GTN 00305970375234
SN 12345678901234
EXP MMMYYYY
LOT 1234567

10010941-01

NDC 0597-0375-23

CYLTEZO® PEN
(adalimumab-adbm)
Injection



ATTENTION PHARMACIST:
Each Patient is Required to Receive the Enclosed Medication Guide.

The Entire Carton is to be Dispensed as a Unit.

Needle Cap for Pen Contains Dry Natural Rubber.

40 mg/0.8 mL

For Subcutaneous Use Only

Rx only



4 Single-Dose Prefilled Pens

Starter Package for Psoriasis



CYLTEZO® PEN
(adalimumab-adbm)
Injection





40 mg/0.8 mL

For Subcutaneous Use Only


4 Single-Dose Prefilled Pens

Starter Package for Psoriasis

QUICKTIPS for USING CYLTEZO® Pen (adalimumab-adbm) Injection
For complete instructions refer to the Instructions for Use in the carton.
Do not try to inject CYLTEZO® yourself until you have been shown the right way to give the injection.

1 Choose Injection Site and Clean	2 Inspect and Uncap	3 Prepare Injection	4 Administer Dose
 <p>Abdomen or Thighs</p>		 <p>Squeeze around injection site</p> <p>Press and hold 90°</p> <p>Tip</p>	 <p>Press button 1 time and slowly count to 10</p> <p>Plunger</p>
<ul style="list-style-type: none">Choose an area on your upper thighs or belly, except for an area 2 inches around your belly button.Choose a different injection site each time you inject, at least 1 inch away from the previous injection site.Clean the injection site with an alcohol wipe.	<ul style="list-style-type: none">Inspect the pen, medicine, and expiration date.Pull the cap straight off the pen.Do Not twist the cap.	<ul style="list-style-type: none">Gently squeeze the skin around the injection site and hold firmly.Hold the pen straight up from the injection site (at a 90-degree angle).Press and hold the tip firmly against squeezed skin at the injection site. The injection button will unlock and be ready for use.	<ul style="list-style-type: none">Press the injection button 1 time. The 'click' sound indicates injection start.Keep holding the pen against the squeezed skin around the injection site while you slowly count to 10.Do Not move the pen during injection.You will see the plunger in the window at the bottom of the pen when the injection is complete.

For more information about CYLTEZO®, call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.



NDC 0597-0375-16

CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

6 Single-Dose Prefilled Pens

Citrate Free

Product and trademark licensed from:
Boehringer Ingelheim International GmbH
© 2023 Boehringer Ingelheim International GmbH
ALL RIGHTS RESERVED
Manufactured by:
Boehringer Ingelheim Pharmaceuticals, Inc.
Ridgefield, CT 06377 USA
Product of USA
US License Number 2006
C77419D

Starter Package for Crohn's Disease, Ulcerative Colitis
or Hidradenitis Suppurativa

Do not use beyond the expiration date.



Starter Package for Crohn's Disease, Ulcerative Colitis
or Hidradenitis Suppurativa

For Subcutaneous Use Only

This Carton Contains:

- 6 dose trays (each containing 1 single-dose prefilled pen with a fixed 1/4 inch length needle)
- 8 alcohol preps
- 1 Package Insert
- 1 Medication Guide
- 1 Instructions for Use

Recommended Dosage: See Prescribing Information.
Needle Cap for Pen Contains Dry Natural Rubber.

Each Sterile Single-Dose Prefilled Pen Contains:
Adalimumab-adbm.....40 mg
Glacial acetic acid.....0.13 mg
Polysorbate 80.....0.8 mg
Sodium acetate trihydrate.....2.4 mg
Trehalose dihydrate.....55.0 mg
Water for Injection, USP

Contains no preservatives.

Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light. **DO NOT FREEZE.** If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use. Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days. Write the date removed from the refrigerator _____

Starter Package for Crohn's Disease, Ulcerative Colitis
or Hidradenitis Suppurativa

40 mg/0.8 mL

For Subcutaneous Use Only

6 Single-Dose Prefilled Pens

For more information about CYLTEZO, including current prescribing information and Medication Guide, go to www.cyltezo.com, scan the code, or call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.



CYLTEZO® PEN
(adalimumab-adbm)
Injection

NDC 0597-0375-16

CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL

NDC 0597-0375-16

NOTE TO PHARMACIST:
(Place Prescription Label Here)

6 Single-Dose Prefilled Pens

For Subcutaneous Use Only

40 mg/0.8 mL

CYLTEZO® PEN
(adalimumab-adbm)
Injection



GTIN 00305970375166
SN 12345678901234
EXP MMYYYY
LOT 1234567

10-7601001

NDC 0597-0375-16

ATTENTION PHARMACIST:
Each Patient is Required to Receive the
Enclosed Medication Guide.

The Entire Carton is to be Dispensed as a Unit.
Needle Cap for Pen Contains Dry Natural Rubber.

CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

Rx only



6 Single-Dose Prefilled Pens

Starter Package for Crohn's Disease, Ulcerative Colitis
or Hidradenitis Suppurativa

Boehringer
Ingelheim

CYLTEZO® PEN
(adalimumab-adbm)
Injection





40 mg/0.8 mL

For Subcutaneous Use Only


6 Single-Dose Prefilled Pens

Starter Package for Crohn's Disease, Ulcerative Colitis
or Hidradenitis Suppurativa

QUICKTIPS for USING CYLTEZO® Pen (adalimumab-adbm) Injection
For complete instructions refer to the Instructions for Use in the carton.
Do not try to inject CYLTEZO® yourself until you have been shown the right way to give the injection.

1 Choose Injection Site and Clean	2 Inspect and Uncap	3 Prepare Injection	4 Administer Dose
 <p>Abdomen or Thighs</p>		 <p>Squeeze around injection site</p> <p>Press and hold 90°</p> <p>Tip</p>	 <p>Press button 1 time and slowly count to 10</p> <p>Plunger</p>
<ul style="list-style-type: none">• Choose an area on your upper thighs or belly, except for an area 2 inches around your belly button.• Choose a different injection site each time you inject, at least 1 inch away from the previous injection site.• Clean the injection site with an alcohol wipe.	<ul style="list-style-type: none">• Inspect the pen, medicine, and expiration date.• Pull the cap straight off the pen.• Do Not twist the cap.	<ul style="list-style-type: none">• Gently squeeze the skin around the injection site and hold firmly.• Hold the pen straight up from the injection site (at a 90-degree angle).• Press and hold the tip firmly against squeezed skin at the injection site. The injection button will unlock and be ready for use.	<ul style="list-style-type: none">• Press the injection button 1 time. The 'click' sound indicates injection start.• Keep holding the pen against the squeezed skin around the injection site while you slowly count to 10.• Do Not move the pen during injection.• You will see the plunger in the window at the bottom of the pen when the injection is complete.

For more information about CYLTEZO®, call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.



Do not use beyond the expiration date.



10010944-04

NDC 0597-0370-82

CYLTEZO[®]
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

2 Single-Dose Prefilled Syringes

Citrate Free

Product and trademark licensed from:
Boehringer Ingelheim International GmbH
© 2022 Boehringer Ingelheim International GmbH
ALL RIGHTS RESERVED
Manufactured by:
Boehringer Ingelheim Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
Product of the USA
US License Number 2006
CT7422H

For Subcutaneous Use Only

This Carton Contains:
• 2 dose trays (each containing 1 single-dose prefilled syringe with a fixed 1/2 inch length needle)
• 2 alcohol preps
• 1 Package Insert
• 1 Medication Guide
• 1 Instructions for Use

Recommended Dosage: See Prescribing Information.
Needle Cap for Pen Contains Dry Natural Rubber.



(01)00305970370826

Each Sterile Single-Dose Prefilled Syringe Contains:
Adalimumab-adbm.....40 mg
Glacial acetic acid.....0.13 mg
Polysorbate 80.....0.8 mg
Sodium acetate trihydrate.....2.4 mg
Trehalose dihydrate.....65.0 mg
Water for Injection, USP

Contains no preservatives.
Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light. **DO NOT FREEZE.** If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use. Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days.

Write the date removed from the refrigerator _____ / ____ / ____.

40 mg/0.8 mL

CYLTEZO[®]
(adalimumab-adbm)
Injection

NDC 0597-0370-82

NOTE TO PHARMACIST:
(Place Prescription Label Here)

2 Single-Dose Prefilled Syringes

40 mg/0.8 mL

For Subcutaneous Use Only

CYLTEZO[®]
(adalimumab-adbm)
Injection

NDC 0597-0370-82

For more information about CYLTEZO, including current prescribing information and Medication Guide, go to www.cyltezo.com, scan the code, or call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.



2 Single-Dose Prefilled Syringes

For Subcutaneous Use Only

40 mg/0.8 mL

CYLTEZO[®]
(adalimumab-adbm)
Injection

NDC 0597-0370-82



GTIN 00305970370826
SN 12345678901234
EXP MMYYYY
LOT 1234567

90-779601001

NDC 0597-0370-82

CYLTEZO[®]
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

R_x only

2 Single-Dose Prefilled Syringes

ATTENTION PHARMACIST:
Each Patient is Required to Receive the Enclosed Medication Guide.
The Entire Carton is to be Dispensed as a Unit.
Needle Cap for Syringe Contains Dry Natural Rubber.



Boehringer
Ingelheim





CYLTEZO[®]
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

2 Single-Dose Prefilled Syringes

QUICKTIPS for USING CYLTEZO® (adalimumab-adbm) Injection
For complete instructions refer to the Instructions for Use in the carton.
Do not try to inject CYLTEZO® yourself until you have been shown the right way to give the injection.

1 Choose Injection Site and Clean	2 Inspect and Uncap	3 Prepare Injection	4 Administer Dose
 <p>Abdomen or Thighs</p>		 <p>Squeeze around injection site</p> <p>Insert needle</p> <p>45°</p>	 <p>Plunger</p>
<ul style="list-style-type: none">• Choose an area on your upper thighs or belly, except for an area 2 inches around your belly button.• Choose a different injection site each time you inject, at least 1 inch away from the previous injection site.• Clean the injection site with an alcohol wipe.	<ul style="list-style-type: none">• Inspect the prefilled syringe, medicine, and expiration date.• Pull the cap straight off the syringe. Do Not let the needle touch anything.• Do Not recap the needle.	<ul style="list-style-type: none">• Squeeze the skin around the injection site and hold firmly.• Hold the syringe at about a 45 degree angle to the injection site.• Insert the needle into the injection site with a quick, smooth motion.	<ul style="list-style-type: none">• Slowly push the plunger down fully until the plunger reaches bottom of syringe.• Remove the needle from the injection site at the same angle you inserted it (45 degree angle).

For more information about CYLTEZO®, call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.



Do not use beyond the expiration date.

NDC 0597-0405-80
CYLTEZO[®]
(adalimumab-adbm)
Injection
20 mg/0.4 mL
For Subcutaneous Use Only
2 Single-Dose Prefilled Syringes

Citrate Free

Product and trademark licensed from:
Boehringer Ingelheim International GmbH
© 2022 Boehringer Ingelheim International GmbH
ALL RIGHTS RESERVED
Manufactured by:
Boehringer Ingelheim Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
Product of the USA
US License Number 2006
CT7974G

This Carton Contains:
• 2 dose trays (each containing 1 single-dose prefilled syringe with a fixed 1/2 inch length needle)
• 2 alcohol preps
• 1 Package Insert
• 1 Medication Guide
• 1 Instructions for Use
Recommended Dosage: See Prescribing Information.
Needle Cap for Pen Contains Dry Natural Rubber.

For Subcutaneous Use Only
Each Sterile Single-Dose Prefilled Syringe Contains:
Adalimumab-adbm.....20 mg
Citric acid.....0.06 mg
Polysorbate 80.....0.4 mg
Sodium acetate trihydrate.....1.21 mg
Trehalose dihydrate.....32.5 mg
Water for Injection, USP
Contains no preservatives.
Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light. **DO NOT FREEZE.** If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use. Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days.
Write the date removed from the refrigerator _____

For more information about CYLTEZO, including current prescribing information and Medication Guide, go to www.cyltezo.com, scan the code, or call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.




NDC 0597-0405-80
CYLTEZO[®]
(adalimumab-adbm)
Injection
20 mg/0.4 mL
For Subcutaneous Use Only
2 Single-Dose Prefilled Syringes


NOTE TO PHARMACIST:
(Place Prescription Label Here)

CYLTEZO[®]
(adalimumab-adbm)
Injection
NDC 0597-0405-80
2 Single-Dose Prefilled Syringes
For Subcutaneous Use Only

NDC 0597-0405-80
CYLTEZO[®]
(adalimumab-adbm)
Injection
20 mg/0.4 mL
For Subcutaneous Use Only



GTN 00305970405801
SN 12345678901234
EXP MMMYYY
LOT 1234567



NDC 0597-0405-80
CYLTEZO[®]
(adalimumab-adbm)
Injection

ATTENTION PHARMACIST:
Each Patient is Required to Receive the Enclosed Medication Guide.
The Entire Carton is to be Dispensed as a Unit.
Needle Cap for Syringe Contains Dry Natural Rubber.

20 mg/0.4 mL

For Subcutaneous Use Only
Rx only



2 Single-Dose Prefilled Syringes







CYLTEZO[®]
(adalimumab-adbm)
Injection

20 mg/0.4 mL

For Subcutaneous Use Only 2 Single-Dose Prefilled Syringes

QUICKTIPS for USING CYLTEZO® (adalimumab-adbm) Injection
For complete instructions refer to the Instructions for Use in the carton.
Do not try to inject CYLTEZO® yourself until you have been shown the right way to give the injection.

1 Choose Injection Site and Clean	2 Inspect and Uncap	3 Prepare Injection	4 Administer Dose
 <p>Abdomen or Thighs</p>		 <p>Squeeze around injection site</p> <p>Insert needle</p> <p>45°</p>	 <p>Plunger</p>
<ul style="list-style-type: none">• Choose an area on your upper thighs or belly, except for an area 2 inches around your belly button.• Choose a different injection site each time you inject, at least 1 inch away from the previous injection site.• Clean the injection site with an alcohol wipe.	<ul style="list-style-type: none">• Inspect the prefilled syringe, medicine, and expiration date.• Pull the cap straight off the syringe. Do Not let the needle touch anything.• Do Not recap the needle.	<ul style="list-style-type: none">• Squeeze the skin around the injection site and hold firmly.• Hold the syringe at about a 45 degree angle to the injection site.• Insert the needle into the injection site with a quick, smooth motion.	<ul style="list-style-type: none">• Slowly push the plunger down fully until the plunger reaches bottom of syringe.• Remove the needle from the injection site at the same angle you inserted it (45 degree angle).

For more information about CYLTEZO®, call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.

CUTSCORE

CUTSCORE



Do not use beyond the expiration date.

NDC 0597-0370-72

CYLTEZO[®]
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

1 Single-Dose Prefilled Syringe

Product and trademark licensed from:
Boehringer Ingelheim
International GmbH

© 2022 Boehringer Ingelheim
International GmbH
ALL RIGHTS RESERVED

Manufactured by:
Boehringer Ingelheim
Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
Product of the USA
US License Number 2006
CT7594H

Citrate
Free



(01)00305970370727

Needle Cap for Syringe Contains Dry Natural Rubber.
Recommended Dosage: See Prescribing Information.

- 1 Instructions for Use
- 1 Medication Guide
- 1 Package Insert
- 2 Alcohol Preps
- 1 dose tray containing 1 single-dose prefilled syringe with a fixed 1/8 inch length needle

This Carton Contains:

Each Sterile Single-Dose Prefilled Syringe Contains:
Adalimumab-adbm.....40 mg
Glacial acetic acid.....0.3 mg
Polysorbate 80.....2.8 mg
Sodium acetate trihydrate.....2.4 mg
Trehalose dihydrate.....65.0 mg
Water for Injection, USP

Contains no preservatives.

Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light. DO NOT FREEZE. If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use. Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days. Write the date removed from the refrigerator

40 mg/0.8 mL

CYLTEZO[®]
(adalimumab-adbm)
Injection

NDC 0597-0370-72

NOTE TO PHARMACIST:
(Place Label Here)

1 Single-Dose Prefilled Syringe

40 mg/0.8 mL

For Subcutaneous Use Only

CYLTEZO[®]
(adalimumab-adbm)
Injection

NDC 0597-0370-72

For more information about CYLTEZO, including current prescribing information and Medication Guide, go to www.cyltezo.com, scan the code, or call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.



40 mg/0.8 mL

CYLTEZO[®]
(adalimumab-adbm)
Injection

NDC 0597-0370-72

1 Single-Dose Prefilled Syringe

For Subcutaneous Use Only



GTIN 00305970370727
SN 12345678901234
EXP MMYYYY
LOT 1234567

10010943-04

NDC 0597-0370-72

CYLTEZO[®]
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

NOT FOR SALE

Rx only

1 Single-Dose Prefilled Syringe



ATTENTION PHARMACIST:
Each Patient is Required to Receive the
Enclosed Medication Guide.

The Entire Carton is to be Dispensed as a Unit.

Needle Cap for Syringe Contains Dry Natural Rubber.



CYLTEZO[®]
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only





1 Single-Dose Prefilled Syringe

QUICKTIPS for USING CYLTEZO® (adalimumab-adbm) Injection

For complete instructions refer to the Instructions for Use in the carton. Do not try to inject CYLTEZO® yourself until you have been shown the right way to give the injection.

For more information about CYLTEZO®, call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.

CUTSCORE

1 Choose Injection Site and Clean	2 Inspect and Uncap	3 Prepare Injection	4 Administer Dose
			
<ul style="list-style-type: none">• Choose an area on your upper thighs or belly, except for an area 2 inches around your belly button.• Choose a different injection site each time you inject, at least 1 inch away from the previous injection site.• Clean the injection site with an alcohol wipe.	<ul style="list-style-type: none">• Inspect the prefilled syringe, medicine, and expiration date.• Pull the cap straight off the syringe. Do Not let the needle touch anything.• Do Not recap the needle.	<ul style="list-style-type: none">• Squeeze the skin around the injection site and hold firmly.• Hold the syringe at about a 45 degree angle to the injection site.• Insert the needle into the injection site with a quick, smooth motion.	<ul style="list-style-type: none">• Slowly push the plunger down fully until the plunger reaches bottom of syringe.• Remove the needle from the injection site at the same angle you inserted it (45 degree angle).

CUTSCORE

Do not use beyond the expiration date.



NDC 0597-0400-89

CYLTEZO[®]
(adalimumab-adbm)
Injection

10 mg/0.2 mL

For Subcutaneous Use Only

2 Single-Dose Prefilled Syringes

Citrate Free

Product and trademark licensed from:
Boehringer Ingelheim International GmbH
© 2022, Boehringer Ingelheim International GmbH
ALL RIGHTS RESERVED
Manufactured by:
Boehringer Ingelheim Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
Product of the USA
US License Number 2006
CT9994C



Recommended Dosage: See Prescribing Information.
Needle Cap for Pen Contains Dry Natural Rubber.

- 2 dose trays (each containing 1 single-dose prefilled syringe with a fixed 1/2 inch length needle)
- 2 alcohol preps
- 1 Package Insert
- 1 Medication Guide
- 1 Instructions for Use

This Carton Contains:
For Subcutaneous Use Only

Each Sterile Single-Dose Prefilled Syringe Contains:
Adalimumab-adbm.....10 mg
Glacial acetic acid.....0.03 mg
Polysorbate 80.....0.2 mg
Sodium acetate trihydrate.....0.61 mg
Trehalose dihydrate.....16.3 mg
Water for Injection, USP

Contains no preservatives.
Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light. DO NOT FREEZE. If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use. Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days.

Write the date removed from the refrigerator _____

10 mg/0.2 mL

CYLTEZO[®]
(adalimumab-adbm)
Injection

NDC 0597-0400-89

NOTE TO PHARMACIST:
(Place Prescription Label Here)

For more information about CYLTEZO, including current prescribing information and Medication Guide, go to www.cyltezo.com, scan the code, or call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.



CYLTEZO[®]
(adalimumab-adbm)
Injection

10 mg/0.2 mL

For Subcutaneous Use Only

2 Single-Dose Prefilled Syringes

NDC 0597-0400-89

2 Single-Dose Prefilled Syringes

For Subcutaneous Use Only

10 mg/0.2 mL

CYLTEZO[®]
(adalimumab-adbm)
Injection

NDC 0597-0400-89

GTN 00305970400899
SN 12345678901234
EXP MMYYYY
LOT 1234567



10011318-02



NDC 0597-0400-89

CYLTEZO[®]
(adalimumab-adbm)
Injection

10 mg/0.2 mL

For Subcutaneous Use Only
Rx only

2 Single-Dose Prefilled Syringes

ATTENTION PHARMACIST:
Each Patient is Required to Receive the Enclosed Medication Guide.
The Entire Carton is to be Dispensed as a Unit.
Needle Cap for Syringe Contains Dry Natural Rubber.







CYLTEZO[®]
(adalimumab-adbm)
Injection

10 mg/0.2 mL

For Subcutaneous Use Only


2 Single-Dose Prefilled Syringes

QUICKTIPS for USING CYLTEZO® (adalimumab-adbm) Injection
For complete instructions refer to the Instructions for Use in the carton.
Do not try to inject CYLTEZO® yourself until you have been shown the right way to give the injection.

1 Choose Injection Site and Clean	2 Inspect and Uncap	3 Prepare Injection	4 Administer Dose
 <p>Abdomen or Thighs</p>		 <p>Squeeze around injection site</p> <p>Insert needle</p> <p>45°</p>	 <p>Plunger</p>
<ul style="list-style-type: none">• Choose an area on your upper thighs or belly, except for an area 2 inches around your belly button.• Choose a different injection site each time you inject, at least 1 inch away from the previous injection site.• Clean the injection site with an alcohol wipe.	<ul style="list-style-type: none">• Inspect the prefilled syringe, medicine, and expiration date.• Pull the cap straight off the syringe. Do Not let the needle touch anything.• Do Not recap the needle.	<ul style="list-style-type: none">• Squeeze the skin around the injection site and hold firmly.• Hold the syringe at about a 45 degree angle to the injection site.• Insert the needle into the injection site with a quick, smooth motion.	<ul style="list-style-type: none">• Slowly push the plunger down fully until the plunger reaches bottom of syringe.• Remove the needle from the injection site at the same angle you inserted it (45 degree angle).

For more information about CYLTEZO®, call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.


10011323-01



Do not use beyond the expiration date.

NDC 0597-0375-48

For more information about CYLTEZO, including current prescribing information and Medication Guide, go to www.cyltezo.com, scan the code, or call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.




CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL
For Subcutaneous Use Only

1 Single-Dose Prefilled Pen

10011323-01



EXP
LOT
MMMYYY
1234567

CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL
For Subcutaneous Use Only

1 Single-Dose Prefilled Pen


NDC 0597-0375-48

CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only
PROFESSIONAL SAMPLE: NOT FOR SALE
Rx only

1 Single-Dose Prefilled Pen



Boehringer
Ingelheim

CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

1 Single-Dose Prefilled Pen

Store in refrigerator at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light. **DO NOT FREEZE.** If needed, you may store at room temperature to a maximum of 77°F (25°C) for up to 14 days in the original carton until time of use. Once stored at room temperature, do not place back in the refrigerator. Discard after 14 days. Write the date removed from the refrigerator _____ / _____ / _____

Each Sterile Single-Dose Prefilled Pen Contains:
Adalimumab-adbm.....40 mg
Glacial acetic acid.....0.13 mg
Polysorbate 80.....0.8 mg
Sodium acetate trihydrate.....2.4 mg
Trehalose dihydrate.....65.0 mg

Water for Injection, USP

Contains no preservatives.

This Carton Contains:
• 1 dose tray containing 1 single-dose prefilled pen with a fixed ½ inch length needle
• 2 alcohol prep pads
• 1 Package Insert
• 1 Medication Guide
• 1 Instructions for Use

Recommended Dosages: See Prescribing Information.

Needle Cap for Pen Contains Dry Natural Rubber.

40 mg/0.8 mL
Injection
(adalimumab-adbm)

CYLTEZO® PEN

NDC 0597-0375-48

For Subcutaneous Use Only

NDC 0597-0375-48

For Subcutaneous Use Only

1 Single-Dose Prefilled Pen

Each Patient is Required to Receive the Enclosed Medication Guide.

The Entire Carton is to be Dispensed as a Unit.

Needle Cap for Pen Contains Dry Natural Rubber.

CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only

1 Single-Dose Prefilled Pen

NDC 0597-0375-48

CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL
For Subcutaneous Use Only

1 Single-Dose Prefilled Pen

Citrate Free

Product and trademark licensed from:
Boehringer Ingelheim International GmbH

© 2023 Boehringer Ingelheim International GmbH
ALL RIGHTS RESERVED

Manufactured by:
Boehringer Ingelheim Pharmaceuticals, Inc.
Ridgefield, CT 06877 USA
Product of the USA
US License Number 2006 CT11334B

NDC 0597-0375-48

For Subcutaneous Use Only

1 Single-Dose Prefilled Pen

CYLTEZO® PEN
(adalimumab-adbm)
Injection

40 mg/0.8 mL

For Subcutaneous Use Only





1 Single-Dose Prefilled Pen

QUICKTIPS for USING CYLTEZO® Pen (adalimumab-adbm) Injection

For complete instructions refer to the Instructions for Use in the carton.

Do not try to inject CYLTEZO® yourself until you have been shown the right way to give the injection.

For more information about CYLTEZO®, call Boehringer Ingelheim Pharmaceuticals, Inc. at 1-800-542-6257.

1 Choose Injection Site and Clean	2 Inspect and Uncap	3 Prepare Injection	4 Administer Dose
 <p>Abdomen or Thighs</p>		 <p>Squeeze around injection site</p> <p>Press and hold 90°</p> <p>Tip</p>	 <p>Press button 1 time and slowly count to 10</p> <p>Plunger</p>
<ul style="list-style-type: none">Choose an area on your upper thighs or belly, except for an area 2 inches around your belly button.Choose a different injection site each time you inject, at least 1 inch away from the previous injection site.Clean the injection site with an alcohol wipe.	<ul style="list-style-type: none">Inspect the pen, medicine, and expiration date.Pull the cap straight off the pen.Do Not twist the cap.	<ul style="list-style-type: none">Gently squeeze the skin around the injection site and hold firmly.Hold the pen straight up from the injection site (at a 90-degree angle).Press and hold the tip firmly against squeezed skin at the injection site. The injection button will unlock and be ready for use.	<ul style="list-style-type: none">Press the injection button 1 time. The 'click' sound indicates injection start.Keep holding the pen against the squeezed skin around the injection site while you slowly count to 10.Do Not move the pen during injection.You will see the plunger in the window at the bottom of the pen when the injection is complete.