




BLA 761074/S-027
 Page 3

<p>No Text Area</p>		<p>NDC 83257-003-01</p> <p>Ogivri[®] (trastuzumab-dkst) For Injection</p> <p>420 mg/vial</p> <p>For intravenous infusion after reconstitution</p> <p>Rx only</p>	<p>BF2254/02</p>
<p>NDC 83257-003-01</p> <p>Biocon Biologics[™]</p> <p>Ogivri[®] (trastuzumab-dkst) For Injection</p> <p>420 mg/vial</p> <p>For intravenous infusion after reconstitution</p> <p>No preservative KEEP REFRIGERATED Multiple-Dose Vial</p> <p>Rx only</p> 	<p>Contents: Each carton contains one 420 mg vial of Ogivri. The content of each Ogivri[®] vial is 420 mg trastuzumab-dkst, D-sorbitol (322.6 mg), L-Histidine (6.0 mg), L-Histidine hydrochloride monohydrate (9.4 mg) and Polyethylene glycol 3350/Macrogol 3350 (94.1 mg). Hydrochloric acid or sodium hydroxide may be used to adjust the pH of the formulation buffer.</p> <p>Store in refrigerator at 2° to 8°C (36° to 46°F) until time of reconstitution.</p> <p>Reconstitution, Dosage, and Administration: Do not use if vacuum is not present. See prescribing information for dosage, preparation, and administration. Reconstitute with 20 mL Bacteriostatic Water for Injection (BWI), USP (0.9% to 1.1% benzyl alcohol) to yield a multiple-dose solution containing 21 mg/mL trastuzumab-dkst that delivers 20 mL (420 mg trastuzumab-dkst). Do Not Shake. Store reconstituted solution in refrigerator at 2° to 8°C (36° to 46°F). Do not Freeze. Discard unused reconstituted solution after 28 days.</p> <p>Do Not Freeze. Do Not Shake Reconstituted Solution. KEEP REFRIGERATED</p>	<p>NDC 83257-003-01</p> <p>Biocon Biologics[™]</p> <p>Ogivri[®] (trastuzumab-dkst) For Injection</p> <p>420 mg/vial</p> <p>For intravenous infusion after reconstitution</p> <p>No preservative KEEP REFRIGERATED Multiple-Dose Vial</p> <p>Rx only</p> 	<p>Biocon Biologics[™]</p>  <p>Manufactured by: Biocon Biologics Inc. 245 Main St, 2nd Floor Cambridge, MA 02142, U.S.A. U.S. License No. 2324</p> <p>Product of India KR/DRUGS/KTK/28D/07/2006</p> <p>Ogivri[®] is a registered trademark of Biosimilars New Co. Ltd.; a Biocon Biologics Company. Copyright © 2023 Biocon Biologics Inc. All rights reserved.</p>

BLA 761074/S-027
 Page 6

NDC 83257-003-11 

Ogivri[®]
 (trastuzumab-dkst)
 For Injection

420 mg/vial

For Intravenous Infusion
 after reconstitution
 Multiple-Dose Vial

Rx only



Store in refrigerator at 2° to 8°C (36° to 46°F) until time of reconstitution.
Reconstitution, Dosage, and Administration: Do not use if vacuum is not present. See prescribing information for dosage, preparation, and administration. Reconstitute with 20 mL of Bacteriostatic Water for Injection (BWF), USP (0.9% to 1.1% benzyl alcohol) to yield a multiple-dose solution containing 21 mg/mL trastuzumab-dkst that delivers 20 mL (420 mg trastuzumab-dkst). Do Not Shake. Store reconstituted solution in refrigerator at 2° to 8°C (36° to 46°F).
 Do not Freeze. Discard unused reconstituted solution after 28 days.

Do Not Freeze.
Do Not Shake Reconstituted Solution.

Manufactured by:
Biocon Biologics Inc.
 245 Main St, 2nd Floor
 Cambridge, MA 02142, U.S.A.
 U.S. License No. 2324
 Product of India
 KR/DRUGS/KTK/28D/07/2006

Date Reconstituted _____/_____/_____
 Discard After _____/_____/_____

Lot: XXXXXXXXX
 Exp.: YYYY-MM

Unvarnish Zone
 40 mmx12mm

BF1815/04




NDC 83257-002-11 

Bacteriostatic Water
 for Injection, USP

NOT FOR USE IN NEWBORNS
 For Drug Diluent Use Only

Multiple-Dose Vial 20 mL

Rx only



Each mL contains: 1.1% benzyl alcohol;
 Water for Injection q.s. pH 4.5 to 7.0.
Use 20 mL as DILUENT for reconstitution of Ogivri[™]
 See enclosed full prescribing information.
 Use only if solution is clear and seal intact.
 Sterile. Non-pyrogenic.
 Store in refrigerator at 2° to 8°C (36° to 46°F).
 Manufactured by: Biocon Biologics Limited,
 Bengaluru, India for Biocon Biologics Inc.,
 Cambridge, MA 02142, U.S.A.

Lot: XXXXXXXXX
 Exp.: YYYY-MM

NVZ
 34.5mm x 10mm

KR/DRUGS/KTK/28/04562019 BF1814/04